arXiv:0708.3389v2 [math.DG] 31 Aug 2007

On the almost sure spiraling of geodesics in negatively curved manifolds

Sa'ar Hersonsky Frédéric Paulin

February 5, 2008

Abstract

Given a negatively curved geodesic metric space M, we study the statistical asymptotic penetration behavior of (locally) geodesic lines of M in small neighborhoods of points, of closed geodesics, and of other compact (locally) convex subsets of M. We prove Khintchine-type and logarithm law-type results for the spiraling of geodesic lines around these objets. As a consequence in the tree setting, we obtain Diophantine approximation results of elements of non-archimedian local fields by quadratic irrational ones. [1]

1 Introduction

Let M be a compact connected Riemannian manifold with negative sectional curvature. Endow the total space of the unit tangent bundle $\pi : T^1M \rightarrow M$ with the Bowen-Margulis measure μ, which is the maximal entropy probability measure for the geodesic flow $(\phi_t)_{t\in\mathbb{R}}$ on T^1M. Let h be the topological entropy of $(\phi_t)_{t\in\mathbb{R}}$. In this paper, we study the statistical asymptotic penetration behavior of (locally) geodesic lines in various objets in M, as tubular neighborhoods of closed geodesic, tubular neighborhoods of compact embedded totally geodesic submanifolds, and other convex subsets. In this introduction, we fix a Lipschitz map $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$.

We first consider a closed geodesic C in M, and study the spiraling of geodesics around C. As the geodesic flow is ergodic with respect to μ, almost every orbit in T^1M is dense. Two geodesic lines, having at some time their unit tangent vectors close, follow themselves closely a long time. Hence almost every geodesic line will stay for arbitrarily long periods of times in a given small tubular neighborhood of C. In this paper, we make this behaviour quantitative. For that, we prove a Khintchine-type theorem, and a logarithm law-type corollary, for geodesic lines spiraling around C. Fix a small enough $\epsilon > 0$, and let $\mathcal{N}_\epsilon C$ be the (closed) ϵ-neighborhood of C.

Theorem 1.1 *If $\int_1^{+\infty} e^{-h\,g(t)}\,dt$ converges (resp. diverges), then for μ-almost no (resp. every) $v \in T^1M$, there exist positive times $(t_n)_{n\in\mathbb{N}}$ converging to $+\infty$ such that $\pi \circ \phi_t(v)$ belongs to $\mathcal{N}_\epsilon C$ for every t in $[t_n, t_n + g(t_n)]$.*

---

[1] **Keywords:** geodesic flow, negative curvature, spiraling, Khintchine theorem, logarithm law, Diophantine approximation, quadratic irrational. **AMS codes:** 53 C 22, 37 D 40, 11 J 61, 30 F 40, 37 A 45, 11 J 83

Define the penetration map $\mathfrak{p} : T^1M \times \mathbb{R} \to [0, +\infty]$ in $\mathcal{N}_\epsilon C$ by $\mathfrak{p}(v,t) = 0$ if $\pi \circ \phi_t(v) \notin \mathcal{N}_\epsilon C$, and otherwise $\mathfrak{p}(v,t)$ is the maximal length of an interval I in $\mathbb{R}$ containing t such that $\pi \circ \phi_s(v) \in \mathcal{N}_\epsilon C$ for every s in I. We refer to [PP] for (many) other ways to measure the penetration of a geodesic line in the ϵ-neighborhood of C.

Corollary 1.2 *For μ-almost every $v \in T^1M$,*

$$\limsup_{t\to+\infty} \frac{\mathfrak{p}(v,t)}{\log t} = \frac{1}{h}\ .$$

When M has constant curvature, and after a geometric translation, Theorem 1.1 and Corollary 1.2 follow from known results (see for instance [DMPV], as well as the recent [BV], where the methods are very different).

We also prove a Khintchine-type theorem for geodesic lines spiraling around totally geodesic submanifolds. For the sake of simplicity in this introduction, we only formulate it for real hyperbolic manifolds, see Theorem 4.9 for a more general statement.

Theorem 1.3 *Assume furthermore that M is a real hyperbolic n-manifold, and C a closed embedded totally geodesic submanifold of dimension $k \geq 1$. Let $\epsilon > 0$ be small enough.*

If $\int_1^{+\infty} e^{-(n-k)g(t)}\ dt$ converges (resp. diverges), then for μ-almost no (resp. every) $v \in T^1M$, there exist positive times $(t_n)_{n\in\mathbb{N}}$ converging to $+\infty$ such that $\pi \circ \phi_t(v)$ belongs to $\mathcal{N}_\epsilon C$ for every t in $[t_n, t_n + g(t_n)]$.

Besides totally geodesic submanifolds, one could also measure the asymptotic spiraling of geodesic lines around other convex subsets, in particular in hyperbolic 3-manifolds. Recall that a subgroup Γ_0 of a Kleinian group Γ, whose limit set is denoted by $\Lambda\Gamma_0$, is *precisely invariant* if the intersection $\gamma\Lambda\Gamma_0 \cap \Lambda\Gamma_0$ is empty for every $\gamma \in \Gamma - \Gamma_0$.

Theorem 1.4 *Assume furthermore that $M = \Gamma\backslash\mathbb{H}^3_\mathbb{R}$ is an hyperbolic 3-manifold, and let Γ_0 be a precisely invariant quasi-fuschian subgroup of Γ. Let δ_0 be the Hausdorff dimension of the limit set $\Lambda\Gamma_0$, and let C_0 be the image in M of the convex hull of $\Lambda\Gamma_0$ in $\mathbb{H}^3_\mathbb{R}$. Let $\epsilon > 0$ be small enough.*

If $\int_1^{+\infty} e^{-(2-\delta_0)g(t)}\ dt$ converges (resp. diverges), then for μ-almost no (resp. every) $v \in T^1M$, there exist positive times $(t_n)_{n\in\mathbb{N}}$ converging to $+\infty$ such that $\pi \circ \phi_t(v)$ belongs to $\mathcal{N}_\epsilon C_0$ for every t in $[t_n, t_n + g(t_n)]$.

All these results follow from our main result, Theorem 4.9, which is much stronger than the above ones. We do not require M to be compact. The first two statements above remain valid when M is complete, non elementary, with a negative upper bound on its sectional curvature, up to replacing h by the critical exponent δ of the fundamental group Γ of M acting on a universal Riemannian covering $\widetilde{M}$ of M, and under some assumptions on Γ. Under these hypotheses on M, Theorem 1.4 is still valid, up to replacing $\mathbb{H}^3_\mathbb{R}$ by $\widetilde{M}$, Γ_0 by a malnormal infinite index convex-cocompact subgroup of Γ with crititical exponent δ_0, and $2 - \delta_0$ by $\delta - \delta_0$.

Furthermore, we do not need $\widetilde{M}$ to be a manifold, Theorem 4.9 is valid in general CAT(-1) spaces, for instance in hyperbolic buildings (i.e. Tits buildings modeled on an hyperbolic reflection group, see [Bou2, GP, HaP] for examples). Corollary 4.11 is an example of an application. In this introduction, we quote two results in the setting of trees. The first one will be proved in Section 4. Let $E[\cdot]$ be the integer part map.

Proposition 1.5 *Let T be a locally finite tree, and $\mathrm{Aut}(T)$ be its locally compact group of automorphisms, such that $\mathrm{Aut}(T)\backslash T$ is finite. Let Γ be a lattice in $\mathrm{Aut}(T)$ acting without inversion on T. Let $\Gamma\backslash\mathcal{G}T$ be the quotient by Γ of the space $\mathcal{G}T$ of isometric maps $\ell : \mathbb{R} \to T$, endowed with its geodesic flow $(\phi_t)_{t\in\mathbb{R}}$ (the action of $\mathbb{R}$ by translations at the source). Let μ be the maximal entropy measure for the action of $(\phi_t)_{t\in\mathbb{R}}$ on $\Gamma\backslash\mathcal{G}T$, and h be its topological entropy. Let C be a cycle in the graph $\Gamma\backslash T$ with L edges.*

If $\int_1^{+\infty} e^{-\frac{h}{L}g(t)}\,dt$ converges (resp. diverges), then for μ-almost no (resp. every) $\ell \in \Gamma\backslash\mathcal{G}T$, there exist positive times $(t_n)_{n\in\mathbb{N}}$ converging to $+\infty$ such that the path $t \mapsto \ell(t)$, starting at time t_n, turns around C at least $E[g(t_n)]$ times.

The next result (see Section 5) uses the Bruhat-Tits tree of the algebraic group SL_2 over the local field $\widehat{K} = \mathbb{F}_q((X))$ of formal Laurent series in the variable X^{-1} over the finite field $\mathbb{F}_q$. Let μ be a Haar measure of $\widehat{K}$. Let $|\cdot|_\infty$ be the absolute value of $\widehat{K}$. Recall that an element of $\widehat{K}$ is *irrational* if it does not belong to the subfield $\mathbb{F}_q(X)$ of rational fractions over $\mathbb{F}_q$, and is *quadratic* if it is a solution of a quadratic equation with coefficients in $\mathbb{F}_q(X)$. The group $\mathrm{SL}_2(\mathbb{F}_q[X])$ acts by homographies on the set of quadratic irrational elements of $\widehat{K}$, and two of these are *congruent* if they are in the same orbit. For every irrational quadratic element α in $\widehat{K}$, let $h(\alpha) = |\alpha - \alpha^*|_\infty^{-1}$, where α^* is the Galois conjugate of α, be its height (see Section 5). In [HP5], we proved a 0-1 measure result for the Diophantine approximation of elements of $\widehat{K}$ by rational ones. The following result (see Section 5) is an analogous one for the Diophantine approximation of elements of $\widehat{K}$ by quadratic irrational ones.

Theorem 1.6 *Let $\varphi : \mathbb{R}_+ \to \mathbb{R}_+^*$ be a map with $u \mapsto \log\varphi(e^u)$ Lipschitz. If the integral $\int_1^{+\infty} \varphi(t)/t\,dt$ diverges (resp. converges), then for μ-almost every $x \in \widehat{K}$,*

$$\liminf \frac{h(\beta)}{\varphi(h(\beta))}\,|x-\beta|_\infty = 0 \text{ (resp. } = +\infty) ,$$

where the lower limit is taken over the quadratic irrational elements β in $\widehat{K}$, in any (resp. some) congruence class, with $h(\beta) \to +\infty$.

For other number theoretic applications of the results of this paper and of [PP], we refer to [HPP].

We first start in Section 2 by recalling properties of the CAT(-1)-spaces X and their spaces at infinity $\partial_\infty X$. We introduce, for every non empty closed convex subset C of X, a nice new distance-like map d_C on $\partial_\infty X - \partial_\infty C$. It generalizes Gromov's distance when C is reduced to a point (see for instance [Bou1]), or Hamenstädt's distance when C is a horoball (see [Ham][HP1, Appendix]).

In Section 3, we present the main technical tool of this paper, a geometric avatar of the Borel-Cantelli lemma. This tool will also be used in Section 6 (which can be read independently of Section 4) to prove other approximation results, both of Khintchine-type and logarithm law-type, expressing how exactly close to a given point passes almost every geodesic line. The following result is joint work with C. S. Aravinda. See Theorem 6.3 for a more general statement about the approximation of given points by almost every geodesic lines.

Theorem 1.7 *Assume furthermore that M has constant curvature -1 and dimension n, and let $x_0 \in M$. Then for Liouville-almost every v in T^1M,*

$$\liminf_{t\to+\infty} \;\frac{d(\pi\circ\phi_t(v), x_0)}{\log t} \;=\frac{1}{n-1}\;.$$

In Section 4, we start by explaining the general situation covering all results 1.1-1.6. We prove some new estimates on the relative geometry of the convex hulls of subgroups of a discrete group of isometries of a CAT(-1) metric space. Among the new geometric informations (possibly useful for later applications), we give in Theorem 4.4 a fluctuating density result. It explains the variation in ϵ of the mass for a conformal density of the ϵ-neighborhood of the limit set of a convex-cocompact subgroup. We then prove our main result, Theorem 4.9.

Khintchine-type theorems and their logarithm law-type corollaries for the spiraling of geodesic lines around cusps were obtained by D. Sullivan [Sul] when M has finite volume and constant curvature, by D. Kleinbock and G. Margulis [KM] if M is a finite volume locally symmetric space, by B. Stratmann and S. L. Velani [SV] (see also for instance [DMPV, BV]) if M is geometrically finite with constant curvature, and by the authors [HP4] if M is geometrically finite with variable curvature. In this paper, which requires many new geometric inputs, our intellectual debt to D. Sullivan's work is still important.

Acknowledgements. The second author acknowledges the support of the University of Georgia at Athens for fruitful visits. Theorem 6.3, and most of the content of Section 6, was essentially proved by Aravinda and the second author during a stay of Aravinda at the Université d'Orsay in 2000. We also thank V. Kleptsyn and M. Pollicot for their comments, and F. Haglund for Remark 4.10.

2 On convexity properties of CAT(-1)-spaces and their discrete subgroups

We refer for instance to [Bou1, BH] for the definitions and basic properties of CAT(-1) metric spaces, their horospheres and their discrete groups of isometries. The new result in this section is the contruction of the distance-like map d_C for a convex subset C.

Let X be a proper CAT(-1) geodesic metric space. Its boundary at infinity is denoted by $\partial_\infty X$. The space of geodesic lines $\ell : \mathbb{R} \to X$ in X, with the compact-open topology, is denoted by $\mathcal{G}X$. The geodesic flow on $\mathcal{G}X$ is the action of $\mathbb{R}$ by translation at the source. For every point x_0 in X, the space $\mathcal{G}X$ identifies with $((\partial_\infty X \times \partial_\infty X) - \Delta) \times \mathbb{R}$, where Δ is the diagonal in $\partial_\infty X \times \partial_\infty X$, by the map which associates to a geodesic lines ℓ the triple $(\ell(-\infty), \ell(+\infty), t)$ of the points at infinity of ℓ and the algebraic distance t on ℓ (oriented from $\ell(-\infty)$ to $\ell(+\infty)$) between $\ell(0)$ and the closest point of ℓ to x_0. Given another base point x'_0, this parametrization (called Hopf's) differs from the one defined by x'_0 only by an additive constant on the third factor. When X is a Riemannian manifold, the map $T^1X \to \mathcal{G}X$ which associates to $v \in T^1X$ the geodesic line $t \mapsto \pi \circ \phi_t(v)$, with $\pi : T^1X \to X$ the canonical projection, is an homeomorphism equivariant with respect to the actions of the isometry group of X and of the geodesic flows on T^1X and $\mathcal{G}X$.

For every ξ in $\partial_\infty X$, the *Buseman function at* ξ is the map β_ξ from $X \times X$ to $\mathbb{R}$ defined by

$$\beta_\xi(x, y) = \lim_{t \to +\infty} d(x, \xi_t) - d(y, \xi_t) ,$$

for any geodesic ray $t \mapsto \xi_t$ ending at ξ. For every subset A of X and every point x in $X \cup \partial_\infty X$, we denote by ${}_xA$ the *shadow of* A *seen from* x, i.e. the set of points at infinity of the geodesic rays or lines starting from x and meeting A.

The following result, needed only in Section 6, is Proposition 3.1 of [HP2] (whose proof of the left inclusion is valid under the only assumptions below).

Lemma 2.1 *Let ρ be a geodesic ray in X, with $x = \rho(0)$ and $\xi = \rho(+\infty)$.*

(1) For every $c, t > 0$, the ball $B_{d_x}(\xi, ce^{-t})$ for the visual distance (see Example 1 below) d_x on $\partial_\infty X$ is contained in the shadow ${}_x\big(B(\rho(t), c)\big)$.

(2) If X is a Riemannian manifold with sectional curvature $-a^2 \le K \le -1$, where $a \ge 1$, then there exists $\kappa_1, \kappa_2, \kappa_3 > 0$ such that for every $c \in \,]0, \kappa_3]$ and every $t \ge \kappa_2$, the shadow ${}_x\big(B(\rho(t), c)\big)$ is contained in the ball $B_{d_x}(\xi, \kappa_1\, c^{\frac{1}{a}}\, e^{-t})$. $\square$

Given a point at infinity $\xi \in \partial_\infty X$ and a horosphere H centered at ξ, let $d_{\xi,H}, d'_{\xi,H} : (\partial_\infty X - \{\xi\})^2 \to \mathbb{R}$ be the following maps. Let $\eta, \eta' \in \partial_\infty X - \{\xi\}$. Let $t \mapsto \eta_t$ and $t \mapsto \eta'_t$ be the geodesic lines starting from ξ, passing at time $t = 0$ in H, and converging to η and η', respectively. Define the *Hamenstädt distance* (see [Ham][HP1, Appendix])

$$d_{\xi,H}(\eta, \eta') = \lim_{t \to +\infty} e^{\frac{1}{2} d(\eta_t, \eta'_t) - t} ,$$

which is a distance inducing the original topology on $\partial_\infty X - \{\xi\}$. The *cuspidal distance* (see [HP3]) $d'_{\xi,H}$ is defined as follows: If $\eta \ne \eta'$, then $-\log\big(2\, d'_{\xi,H}(\eta, \eta')\big)$ is the signed distance, along the geodesic line $]\xi, \eta[$ oriented from ξ to η, from H to the horosphere centered at η and meeting the geodesic line $]\xi, \eta'[$ in exactly one point. Though not necessarily an actual distance, $d'_{\xi,H}$ is equivalent to the Hamenstädt distance (see [HP3, Rem. 2.6]).

Let C be non empty closed convex subset of X. (Recall that a subset C in a CAT(-1) metric space is *convex* if C contains the geodesic segment between any two points in C.) We denote by $\partial_\infty C$ its set of points at infinity, and by ∂C its boundary in X. For every ξ in $X \cup \partial_\infty X$, we define *the closest point to* ξ *on the convex set* C, denoted by $\pi_C(\xi)$, to be the following point p in $C \cup \partial_\infty C$: If $\xi \in X$, then p belongs to C and minimizes the distance between x and a point of C; if $\xi \in \partial_\infty X - \partial_\infty C$, then the (closed) horoball centered at ξ whose horosphere contains p meets C exactly at p; if $\xi \in \partial_\infty C$, then we define $p = \xi$. This p exists, is unique, and $\pi_C : X \cup \partial_\infty X \to C \cup \partial_\infty C$ is continuous, by the properties of CAT(-1)-spaces. When X is a Riemannian manifold, π_C is (outside $\partial_\infty C$) the orthogonal projection on C.

Let us define a distance-like map d_C on $\partial_\infty X - \partial_\infty C$. For every ξ, η in $\partial_\infty X - \partial_\infty C$, let $t \mapsto \xi_t$ and $t \mapsto \eta_t$ be geodesic rays, starting at time 0 from the closest points to ξ and η on C, and converging to ξ and η respectively. Define

$$d_C(\xi, \eta) = \lim_{t \to +\infty} e^{\frac{1}{2} d(\xi_t, \eta_t) - t} = \lim_{t \to +\infty} e^{\frac{1}{2}\big(d(\xi_t, \eta_t) - d(\xi_t, \pi_C(\xi)) - d(\eta_t, \pi_C(\eta))\big)} \;.$$

Note that these limits exist, and the second equality holds for every geodesic rays $t \mapsto \xi_t$ and $t \mapsto \eta_t$ converging to ξ and η, respectively. For every isometry γ of X, we have

$$d_{\gamma C}(\gamma \xi, \gamma \eta) = d_C(\xi, \eta) \;.$$

In particular, any isometry of X preserving C preserves d_C. For every $\epsilon > 0$, the (closed) ϵ-neighborhood $\mathcal{N}_\epsilon C$ of C in X is still convex with $\partial_\infty(\mathcal{N}_\epsilon C) = \partial_\infty C$, and

$$d_{\mathcal{N}_\epsilon C}(\xi, \eta) = e^{\epsilon}\; d_C(\xi, \eta) \;.$$

Examples.

(1) If C is reduced to a point x in X, then $d_C = d_x$ is the usual visual distance seen from x on $\partial_\infty X$ (see for instance [Bou1]).

(2) If C is a (closed) ball of center x and radius r, then $d_C = e^r\, d_x$, as $C = \mathcal{N}_r\{x\}$.

(3) If C is a (closed) horoball with point at infinity ξ and boundary horosphere H, then $d_C = d_{\xi,\,H}$ is the Hamenstädt distance on $\partial_\infty X - \{\xi\}$ as recalled above. If $(C_n)_{n \in \mathbb{N}}$ is a sequence of balls converging uniformly on compact subsets of X to an horoball C, then the maps d_{C_n} converge uniformly on compact subsets of $\partial X - \{\xi\}$ to d_C.

(4) If X is a metric tree, then it is easy to prove that, for every ξ, η in $\partial_\infty X - \partial_\infty C$,

$$d_C(\xi, \eta) = \begin{cases} e^{\frac{1}{2}\, d(\pi_C(\xi),\, \pi_C(\eta))} & \text{if } \pi_C(\xi) \neq \pi_C(\eta) \\ e^{-d(\pi_C(\xi),\, q)} & \text{if } \pi_C(\xi) = \pi_C(\eta) \text{ and } [\pi_C(\xi), \xi[\, \cap\, [\pi_C(\eta), \eta[\; = [\pi_C(\xi), q] \;. \end{cases} \tag{-1-}$$

In particular, in a small enough neighborhood of any point ξ_0 in $\partial_\infty X - \partial_\infty C$, the map d_C then coincides with the visual distance $d_{\pi_C(\xi_0)}$.

(5) Let X be the real hyperbolic n-space $\mathbb{H}^n_{\mathbb{R}}$, and let C be a complete totally geodesic submanifold of dimension k with $0 < k < n$. Let x_0 be a point in C, and $S_{x_0}(C^\perp)$ be the sphere of unit tangent vectors at x_0 that are perpendicular to C, endowed with the angular distance $(v, v') \mapsto \angle_{x_0}(v, v')$. Note that the standard Euclidean distance on $S_{x_0}(C^\perp)$ is given by $(v, v') \mapsto 2 \sin \frac{\angle_{x_0}(v, v')}{2}$. For every $\xi \in \partial_\infty X - \partial_\infty C$, let $\pi'_C(\xi)$ be the parallel transport to x_0, along a geodesic line through $x_0, \pi_C(\xi)$, of the unit tangent vector at $\pi_C(\xi)$ of the geodesic ray $[\pi_C(\xi), \xi[$. We thus get a map $\pi'_C : \partial_\infty X - \partial_\infty C \to S_{x_0}(C^\perp)$. In particular, $(\pi_C, \pi'_C) : \partial_\infty X - \partial_\infty C \to C \times S_{x_0}(C^\perp)$ is an homeomorphism.

Lemma 2.2 *For every ξ, η in $\partial_\infty X - \partial_\infty C$,*

$$d_C(\xi, \eta) = \sqrt{\;\sinh^2 \frac{d\big(\pi_C(\xi), \pi_C(\eta)\big)}{2} + \sin^2 \frac{\angle_{x_0}\big(\pi'_C(\xi), \pi'_C(\eta)\big)}{2}\;} \;.$$

Proof. Let $\rho = d(\pi_C(\xi), \pi_C(\eta))$ and let $\theta = \angle_{x_0}(\pi'_C(\xi), \pi'_C(\eta))$. We have to prove that $d_C(\xi, \eta) = \frac{1}{2}\sqrt{e^\rho + e^{-\rho} - 2\cos\theta}$. This last formula follows from an easy computation using the picture below. Recall that $\sinh b = 1/\tan\alpha$, where b is the hyperbolic length of the arc of any half-circle perpendicular to the horizontal plane between the angles α and $\frac{\pi}{2}$ in the upper halfspace model of $\mathbb{H}^3_\mathbb{R}$ (see [Bea, page 145]).

Take a copy of $\mathbb{H}^3_\mathbb{R}$ containing ξ, η and a geodesic line passing through $\pi_C(\xi), \pi_C(\eta)$. Use the upper halfspace model of $\mathbb{H}^3_\mathbb{R}$ where this geodesic line is a vertical line between 0 and ∞, with $\pi_C(\eta)$ above $\pi_C(\xi)$. Scale such that the Euclidean distance between 0 and ξ is 1. Consider the points $\eta'_s, \eta_s, \xi'_s, \xi_s$ at Euclidean height s close to 0 on respectively $[\pi_C(\eta), \eta[\,,\ ]\eta, \xi[$ close to η, $[\pi_C(\xi), \xi[\,,\ ]\eta, \xi[$ close to ξ, so that $d_C(\xi, \eta)$ is equal to

$$\lim_{s \to 0}\ e^{\frac{1}{2}\left(d(\xi_s, \eta_s) - d(\xi'_s, \pi_C(\xi)) - d(\eta'_s, \pi_C(\eta))\right)}\ .$$

Now just use several times the previously mentionned formula $\sinh b = 1/\tan\alpha$. $\square$



In particular, if $X = \mathbb{H}^2_\mathbb{R}$, if C is a geodesic line and if ξ, η are in the same component of $\partial_\infty X - \partial_\infty C$, then

$$d_C(\xi, \eta) = \sinh \frac{d(\pi_C(\xi), \pi_C(\eta))}{2}\ .$$

By taking a, b, c in the same component of $\partial_\infty X - \partial_\infty C$ such that $d(\pi_C(a), \pi_C(b)) = d(\pi_C(b), \pi_C(c)) = \frac{1}{2}d(\pi_C(a), \pi_C(c))$ are big enough, we see that d_C does not satisfy the triangle inequality, hence is not a distance.

After these examples, let us go back to the general situation on X, C, and let us prove some results saying that at least on compact subsets, the map d_C behaves quite like a distance.

Lemma 2.3 *(1) For every x_0 in X, for every compact subset K of $\partial_\infty X - \partial_\infty C$, there exists a constant $c_K > 0$ such that for every ξ, η in K, we have*

$$\frac{1}{c_K} d_{x_0}(\xi, \eta) \leq d_C(\xi, \eta) \leq c_K\, d_{x_0}(\xi, \eta)\ .$$

(2) For every ξ in $\partial_\infty X - \partial_\infty C$, the map $\eta \mapsto d_C(\xi, \eta)$ is proper on $\partial_\infty X - \partial_\infty C$.
(3) For every ξ, η in $\partial_\infty X - \partial_\infty C$,

$$(3 - 2\sqrt{2})\ e^{\frac{1}{2}d(\pi_C(\xi), \pi_C(\eta))}\ e^{-d(C,\,]\xi, \eta[)} \leq d_C(\xi, \eta) \leq e^{\frac{1}{2}d(\pi_C(\xi), \pi_C(\eta))}\ .$$

(4) There exist universal constants $c, c' > 0$ such that for every ξ, η in $\partial_\infty X - \partial_\infty C$, if $d_C(\xi, \eta) \leq c$, then C and the geodesic line $]\xi, \eta[$ are disjoint, and

$$\frac{1}{c'}\ e^{-d(C,\,]\xi, \eta[)}\ \leq d_C(\xi, \eta) \leq\ c'\ e^{-d(C,\,]\xi, \eta[)}\ .$$

Note that by hyperbolicity, $\min\{d(\pi_C(\xi),\pi_C(\eta)), d(C,]\xi,\eta[)\}$ is, for every ξ,η, less than a universal constant.

Proof. For every x_0 in the convex subset C and ξ in $\partial_\infty X-\partial_\infty C$, by the triangle inequality and the CAT(-1) inequality, we have

$$d(\xi_t,x_0)\leq d(\xi_t,\pi_C(\xi))+d(x_0,\pi_C(\xi))\leq d(\xi_t,x_0)+2\log(1+\sqrt{2})\;,$$

with ξ_t as above. Hence for every ξ,η in $\partial_\infty X-\partial_\infty C$,

$$(3-2\sqrt{2})\;d_{x_0}(\xi,\eta)\leq d_C(\xi,\eta)\;e^{-\frac{1}{2}\big(d(x_0,\pi_C(\xi))+d(x_0,\pi_C(\eta))\big)}\leq d_{x_0}(\xi,\eta)\;. \qquad (\text{-}\,2\,\text{-})$$

The first result easily follows. By taking $x_0=\pi_C(\xi)$ in the lower bound of Equation (- 2 -), the second assertion also follows. By taking x_0 to be the midpoint of the geodesic segment $[\pi_C(\xi),\pi_C(\eta)]$ in Equation (- 2 -), and since $d_{x_0}\leq 1$, the upper bound in the third assertion follows.

By the triangle inequality, $d_{x_0}(\xi,\eta)\geq e^{-d(x_0,\,]\xi,\eta[)}$. Hence, by taking x_0 in Equation (- 2 -) to be the closest point of C to $]\xi,\eta[$ if C and $]\xi,\eta[$ are disjoint, or any point in $C\cap]\xi,\eta[$ otherwise, and by using again the triangle inequality, the lower bound in the third assertion follows.

The last assertion follows by standard techniques of approximation by trees (see for example [GH, page 33]). □

In particular, the non negative symmetric map d_C vanishes on and only on the diagonal of $(\partial_\infty X-\partial_\infty C)^2$. But as seen above, d_C is not always a distance.

It also follows from Lemma 2.3 (1) that the uniform structure (see for instance [Bou]) defined (on compacts subsets) by the family $\big(\{(x,y)\in(\partial_\infty X-\partial_\infty C)^2\;:\;d_C(\xi,\eta)\leq\epsilon\}\big)_{\epsilon>0}$ is isomorphic (on compacts subsets) to the uniform structure defined by the distance d_{x_0}.

Remark. Though we won't need it in this paper, here is a formula expressing the distance-like map d_C, when $C=L$ is a geodesic line with endpoints L_-,L_+, in terms of the Hamenstädt distance and the cuspidal distance: For every ξ,η in $\partial_\infty X-\{L_-,L_+\}$, for every horosphere H small enough centered at L_-,

$$d_C(\xi,\eta)=\frac{d_{L_-,H}(\xi,\eta)}{2\big(d'_{L_-,H}(\xi,L_+)d'_{L_-,H}(\eta,L_+)\big)^{\frac{1}{2}}}\;.$$

Proof. Let H_ξ (resp. H_η) be the horosphere centered at ξ (resp. η) passing through $\pi_C(\xi)$ (resp. $\pi_C(\eta)$). Let h_ξ (resp. h_η) be the intersection point of H_ξ (resp. H_η) with the geodesic line $]\xi,L_-[$ (resp. $]\eta,L_-[$). Then

$$d_C(\xi,\eta)=\lim_{t\to+\infty}\;e^{\frac{1}{2}\big(d(\xi_t,\eta_t)-d(\xi_t,h_\xi)-d(\eta_t,h_\eta)\big)}=d_{L_-,H}(\xi,\eta)\;e^{\frac{1}{2}\big(d(h_\xi,H)+d(h_\eta,H)\big)}\;,$$

which proves the result. □

Let Γ be a discrete group of isometries of X. Its limit set is denoted by $\Lambda\Gamma$, and if $\Lambda\Gamma$ contains at least two points, then the convex hull of $\Lambda\Gamma$ is denoted by $\mathcal{C}\Gamma$. Recall that

$\partial_\infty \mathcal{C}\Gamma = \Lambda\Gamma$. The *critical exponent* of Γ is the unique number δ_Γ in $[0, +\infty]$ such that the *Poincaré series* $P_{x_0,\Gamma}(s) = \sum_{\gamma\in\Gamma} e^{-s\, d(x_0,\gamma x_0)}$ of Γ converges for $s > \delta_\Gamma$ and diverges for $s < \delta_\Gamma$, where x_0 is any point in X. The group Γ is called *of divergent type* if its Poincaré series diverges at $s = \delta_\Gamma$. The group Γ is *non elementary* if $\Lambda\Gamma$ contains at least three points, and we have then $\delta_\Gamma > 0$. Note that when X is a Riemannian manifold and Γ is torsion free with compact quotient X/Γ, then the critical exponent δ_Γ of Γ is the topological entropy of the geodesic flow of X/Γ (see for instance [Man]).

If $\delta \in\,]0, +\infty[$, a *conformal* (or Patterson-Sullivan) *density of dimension* δ for Γ is a family $(\mu_x)_{x\in X}$ of finite Borel measures on $\partial_\infty X$, such that

- $\forall\, \gamma \in \Gamma\ ,\ \ \gamma_*\mu_x = \mu_{\gamma x}$,
- $\forall\, x, y \in X,\ \forall\, \xi \in \partial_\infty X\ ,\ \ \frac{d\mu_x}{d\mu_y}(\xi) = e^{-\delta\beta_\xi(x,y)}$.

Using Hopf's parametrization with respect to any base point x_0 of X, the (Patterson-Sullivan-)*Bowen-Margulis measure* associated to this family is the measure $\widetilde{\mu}_{\rm BM}$ on $\mathcal{G}X$ given by

$$d\,\widetilde{\mu}_{\rm BM} = \frac{d\mu_{x_0}(\xi)\, d\mu_{x_0}(\eta)\, dt}{d_{x_0}(\eta,\xi)^{2\delta}}\ .$$

This measure on $\mathcal{G}X$ is independant of x_0, invariant by the action of Γ and by the geodesic flow (and by the time reversal $\ell \mapsto \{t \mapsto \ell(-t)\}$), hence defines a measure $\mu_{\rm BM}$ on $\Gamma\backslash\mathcal{G}X$ which is invariant by the quotient geodesic flow (see for instance [Bou1, Rob]). Note that if $\mu_{\rm BM}$ is finite, then $\delta = \delta_\Gamma$ and Γ is of divergent type (see [Rob, page 18-19]).

If Γ is of divergent type with a finite non zero critical exponent δ, then (see for instance [Bou1]) there exists a conformal density of dimension δ for Γ, which is unique up to a positive scalar factor, and which is ergodic with respect to the action of Γ on $\partial_\infty X$. The Bowen-Margulis measure associated to any such conformal family (both on $\mathcal{G}X$ and on $\Gamma\backslash\mathcal{G}X$) will be called a *Bowen-Margulis measure* of Γ (it is also uniquely defined up to a positive scalar factor). When X is a manifold and Γ acts freely on X with compact quotient, the Bowen-Margulis measure on the unit tangent bundle of the compact negatively curved manifold $M = \Gamma\backslash X$, normalized to be a probability measure, is the maximal entropy probability measure for the geodesic flow of M (via the canonical identification of $\mathcal{G}X$ and T^1X), see for instance [Kai]. When furthermore X has constant curvature, then the Bowen-Margulis measure and the Liouville measure (when both are normalized) coincide on M.

The following result, which is obvious by definition of the Bowen-Margulis measure, will be used in the sections 4 and 6.

Lemma 2.4 *Let $\pi_+ : \mathcal{G}X \to \partial_\infty X$ be the map $\ell \mapsto \ell(+\infty)$. Let $\widetilde{\mu}_{BM}$ be the Bowen-Margulis measure on $\mathcal{G}X$ associated to a conformal family $(\mu_x)_{x\in X}$ for Γ. Then the preimage by π_+ of a set of measure 0 (resp. > 0) for μ_x (for some (equivalently for any) x in X) has measure 0 (resp. > 0) for $\widetilde{\mu}_{BM}$.* □

Besides its invariance under Γ and the geodesic flow, and its ergodicity on $\Gamma\backslash\mathcal{G}X$, this is the only property of the Bowen-Margulis measure $\widetilde{\mu}_{\rm BM}$ on $\mathcal{G}X$ that will be used in this paper. In particular, we may replace $\widetilde{\mu}_{\rm BM}$ by any other measure satisfying these invariance properties and this lemma, as for instance the Knieper measure (see [Kni]).

The group Γ is said to be *convex-cocompact* if $\Lambda\Gamma$ contains at least two points, and if the action of Γ on $\mathcal{C}\Gamma$ has compact quotient. In particular, the group generated by an hyperbolic isometry of X is convex-cocompact, with critical exponent 0. In fact, if Γ is convex-cocompact, then its critical exponent is 0 if and only if Γ has an index two subgroup generated by an hyperbolic isometry of X. Note that $\partial_\infty \mathcal{C}\Gamma = \Lambda\Gamma$, and that if Γ is convex-cocompact then Γ is of divergent type (see for instance [Bou1, Rob]).

For every $f, g : \mathbb{N} \to [0, +\infty[$, write $f \asymp g$ if there exists a constant $c \geq 1$ such that $\frac{1}{c} f \leq g \leq cf$. For every x_0 in X, if Γ is convex-cocompact and non elementary, with critical exponent δ_Γ, then

$$\mathrm{Card}(B(x_0, n) \cap \Gamma x_0) \asymp e^{\delta_\Gamma n}$$

(see for instance [Rob], where others, much more general, assumptions on Γ are given for this property to hold. This is the case for example when the Bowen-Margulis measure $\mu_{\rm BM}$ of Γ is finite (and the length spectrum is non arithmetic), see [Rob, page 56]).

Lemma 2.5 *Let Γ_0 be a convex-cocompact subgroup with infinite index in a discrete group of isometries Γ of X. Let δ_0 and δ be the critical exponents of Γ_0 and Γ respectively. Then $\delta_0 < \delta$.*

Proof. This is well-known (see for instance [Fur] in a special case). □

Recall that the *virtual normalizer* $\mathrm{N}\Gamma_0$ of a convex-cocompact subgroup Γ_0 of Γ is the stabilizer in Γ of the limit set $\Lambda\Gamma_0$. It contains the normalizer of Γ_0 in Γ, and it contains Γ_0 with finite index (see for instance [KS, Arz]).

Recall that a subgroup H of a group G is *malnormal* if, for every g in $G - H$, we have $gHg^{-1} \cap H = \{1\}$. We will say that a subgroup H of a group G is *almost malnormal* if, for every g in $G - H$, the subgroup $gHg^{-1} \cap H$ is finite. Note that malnormal implies almost malnormal, and that the converse is true if the ambient group is torsion free.

The following result is folklore, we provide a proof because we couldn't find a precise reference.

Proposition 2.6 *Let Γ_0 be a convex-cocompact subgroup of a discrete group Γ of isometries of X, then the following assertions are equivalent.*

(1) Γ_0 is almost malnormal in Γ;

(2) the limit set of Γ_0 is precisely invariant, i.e. for every $\gamma \in \Gamma - \Gamma_0$, the set $\Lambda\Gamma_0 \cap \gamma\Lambda\Gamma_0$ is empty;

(3) $\mathcal{C}\Gamma_0 \cap \gamma\mathcal{C}\Gamma_0$ is compact for every $\gamma \in \Gamma - \Gamma_0$;

(4) for every $\epsilon > 0$, there exists $\kappa = \kappa(\epsilon) > 0$ such that $\mathrm{diam}\big(\mathcal{N}_\epsilon \mathcal{C}\Gamma_0 \cap \gamma \mathcal{N}_\epsilon \mathcal{C}\Gamma_0\big) \leq \kappa$ for every $\gamma \in \Gamma - \Gamma_0$.

The convex hull in X of the limit set of a convex-cocompact subgroup is non compact. Hence an almost malnormal convex-cocompact subgroup of Γ is equal to its virtual normalizer, by (3).

Proof. As $\partial_\infty \mathcal{C}\Gamma_0 = \Lambda\Gamma_0$, it is clear that (4) implies (3), which implies (2), which implies (1).

Let us prove that (1) implies (4). Let $C_0 = \mathcal{C}\Gamma_0$ and $\epsilon > 0$. Assume by absurd that for every n in $\mathbb{N}$, there exists γ_n in $\Gamma - \Gamma_0$ and x_n, y_n in $\mathcal{N}_\epsilon C_0 \cap \gamma_n \mathcal{N}_\epsilon C_0$ with $d(x_n, y_n) \geq n$. As $\Gamma_0 \backslash \mathcal{N}_\epsilon C_0$ is compact and the action of Γ_0 is isometric, there exists $R > 0$ such that $\Gamma_0 B(x, R)$ contains $\mathcal{N}_\epsilon C_0$ for every x in $\mathcal{N}_\epsilon C_0$. As γ_n is an isometry, we also have that $\gamma_n \Gamma_0 \gamma_n^{-1} B(y, R)$ contains $\gamma_n \mathcal{N}_\epsilon C_0$, for every y in $\gamma_n \mathcal{N}_\epsilon C_0$. Up to conjugating γ_n by an element of Γ_0, we may assume that x_n stays in a compact subset K of X, and we define $K' = \mathcal{N}_R K$, which is compact. As Γ is discrete, the number N of elements γ in Γ, such that $\gamma K' \cap K'$ is non empty, is finite. As Γ_0 is convex-cocompact, the upper bound of the cardinals of the finite subgroups of Γ_0 is finite. Hence, as Γ_0 is almost malnormal, there exists $N' \in \mathbb{N}$ such that for every γ in $\Gamma - \Gamma_0$, the cardinal of $\gamma \Gamma_0 \gamma^{-1} \cap \Gamma_0$ is at most $N' - 2$. Take n in $\mathbb{N}$ with $n > NN' \operatorname{diam} K'$. Subdivide the segment between x_n and y_n in points $u_0 = x_n, u_1, \dots, u_{NN'} = y_n$, such that $d(u_k, u_{k+1}) > \operatorname{diam} K'$ for $0 \leq k \leq NN' - 1$. As K' contains $B(x_n, R)$ and x_n, y_n belong to the convex subset $\mathcal{N}_\epsilon C_0 \cap \gamma_n \mathcal{N}_\epsilon C_0$, for $0 \leq k \leq NN'$, there exist α_k, β_k in Γ_0 such that $u_k \in \alpha_k K'$ and $u_k \in \gamma_n \beta_k \gamma_n^{-1} K'$. Note that $\alpha_k \neq \alpha_j$ if $k \neq j$, as $d(u_k, u_j) > \operatorname{diam} K'$. By the definition of N, there exists $(k_j)_{1 \leq j \leq N'}$ with $\alpha_{k_j}^{-1} \gamma_n \beta_{k_j} \gamma_n^{-1} = \alpha_{k_1}^{-1} \gamma_n \beta_{k_1} \gamma_n^{-1}$ for $1 \leq j \leq N'$. Hence $\gamma_n \beta_{k_j} \beta_{k_1}^{-1} \gamma_n^{-1} = \alpha_{k_j} \alpha_{k_1}^{-1}$, for $2 \leq j \leq N'$, which contradicts the fact that the cardinal of $\gamma_n \Gamma_0 \gamma_n^{-1} \cap \Gamma_0$ is at most $N' - 2$. □

Remark. The fact that the first two assertions are equivalent follows also from the well-known equality

$$\Lambda\Gamma_0 \cap \gamma\Lambda\Gamma_0 = \Lambda(\Gamma_0 \cap \gamma\Gamma_0\gamma^{-1}) ,$$

see for instance [SS, Coro. 3] for a proof in a special case.

3 A geometric avatar of the Borel-Cantelli lemma

The main technical tool of this paper is the following result, which is a suitable enhancement of the Borel-Cantelli Lemma.

Theorem 3.1 *Let (Z, μ) be a measured space with $\mu(Z)$ finite, and $(B_i(\epsilon))_{i \in I,\, \epsilon \in \,]0, +\infty[}$ a family of measurable subsets in Z, non-decreasing in ϵ (for the inclusion), endowed with a map $i \mapsto n_i$ from I to $\mathbb{N}$ such that $I_n = \{i \in I \;:\; n_i = n\}$ is finite for every n. Let f_1, f_2, f_3, f_4 be maps from $\mathbb{N}$ to $]0, +\infty[$ and f_5 a map from $]0, +\infty[$ to itself. Let E be the (measurable) set of points in Z belonging to infinitely many subsets $B_i(f_3(n_i))$ for i in I.*

[A] Assume that $f_3 \leq f_2$ and that there exists $c \geq 1$ such that, for every n in $\mathbb{N}$, i in I and $\epsilon \in \,]0, f_2(n_i)]$, one has $\operatorname{Card} I_n \leq c f_1(n)$ and $\mu(B_i(\epsilon)) \leq c f_4(n_i) f_5(\epsilon)$. If the series $\sum_{n=0}^{\infty} f_1(n) f_4(n) f_5(f_3(n))$ converges, then $\mu(E) = 0$.

[B] Assume that there exists $c \geq 1$ such that

(1) $f_3 \leq f_2$,

(2) $\frac{1}{f_5 \circ f_2} \leq f_4 f_1$,

(3) there exists $c', c'' > 1$ *such that for every* $\epsilon, \epsilon' > 0$, *if* $\epsilon' \leq c'\epsilon$, *then* $f_5(\epsilon') \leq c'' f_5(\epsilon)$,

(4) for every n *in* $\mathbb{N}$, *one has* $\frac{1}{c} f_1(n) \leq \mathrm{Card}\ I_n \leq c f_1(n)$,

(5) for every i *in* I *and* $\epsilon \in \,]\,0, f_2(n_i)]$, *we have* $\frac{1}{c} f_4(n_i) f_5(\epsilon) \leq \mu(B_i(\epsilon)) \leq c f_4(n_i) f_5(\epsilon)$,

(6) for every n *in* $\mathbb{N}$, *the subsets* $B_i(f_2(n))$ *for* i *in* I_n *are pairwise disjoint,*

(7) for every i, j *in* I *with* $n_i < n_j$, *if the intersection of* $B_j(f_3(n_j))$ *and* $B_i(f_3(n_i))$ *is non empty, then* $B_j(f_2(n_j))$ *is contained in* $B_i(cf_3(n_i))$.

If the series $\sum_{n=0}^{\infty} f_1(n) f_4(n) f_5(f_3(n))$ *diverges, then* $\mu(E) > 0$.

Note that (except for the convergence of the series) every hypothesis of Case [A] is part of an hypothesis (1)-(5) of Case [B]. Hence when checking the hypotheses when we want to apply both cases of this theorem, we will only chek the ones of Case [B].

Proof. For i in I and n in $\mathbb{N}$, let $B_i = B_i(f_3(n_i))$ and $A_n = \bigcup_{i \in I_n} B_i$, so that $E = \bigcap_{n \in \mathbb{N}} \bigcup_{k \geq n} A_k$.

Under the assumptions of [A], by the subadditivity of μ, we have the inequality $\mu(A_n) \leq c^2 f_1(n) f_4(n) f_5(f_3(n))$. Therefore the end of the proof is standard: If the series $\sum_{n=0}^{\infty} f_1(n) f_4(n) f_5(f_3(n))$ converges, then the sequence $u_k = \sum_{n=k}^{\infty} f_1(n) f_4(n) f_5(f_3(n))$ tends to 0, therefore

$$\mu(E) = \lim_{n \to \infty} \mu\left(\bigcup_{k=n}^{\infty} A_k\right) \leq \lim_{n \to \infty} c^2 u_n = 0\ .$$

Assume now that the assumptions of [B] hold. We first claim that

$$f_1(n) f_4(n) f_5(f_3(n)) \leq c^2 \mu(A_n)\ . \quad (*)$$

Indeed, the balls B_i for i in I_n are pairwise disjoint by (1) and (6), since the subsets $B_i(r)$ are non-decreasing in r. By the additivity of μ, by the lower bounds in (4) and (5), the inequality (*) hence follows.

In particular, $\sum \mu(A_n)$ diverges if $\sum f_1(n) f_4(n) f_5(f_3(n))$ diverges.

Now, let n, m be in $\mathbb{N}$ with $n < m$. By the properties (6) and (7), for every i in I_n, we have

$$\mu\left(B_i(cf_3(n_i))\right) \geq \mathrm{Card}\{j \in I_m \ :\ B_j \cap B_i \neq \emptyset\} \min_{j \in I_m} \mu\left(B_j(f_2(m))\right)\ .$$

Hence by (5)

$$\mathrm{Card}\{j \in I_m \ :\ B_j \cap B_i \neq \emptyset\} \leq \frac{c f_4(n) f_5(c f_3(n))}{\frac{1}{c} f_4(m) f_5(f_2(m))}\ . \quad (**)$$

Therefore

$$\begin{aligned}
\mu(A_n \cap A_m) &\leq \textstyle\sum_{i\in I_n} \sum_{j\in I_m\,,\, B_j\cap B_i\neq\emptyset}\ \mu(B_j) \\
&\leq c f_1(n) \times \frac{c f_4(n) f_5(c f_3(n))}{\frac{1}{c} f_4(m) f_5(f_2(m))} \times c f_4(m) f_5(f_3(m)) \\
&\leq c^4 (c'')^{\frac{\log c}{\log c'}+1} f_1(n) f_4(n) f_5(f_3(n)) f_1(m) f_4(m) f_5(f_3(m)) \\
&\leq c^8 (c'')^{\frac{\log c}{\log c'}+1} \mu(A_n)\mu(A_m)\ .
\end{aligned}$$

The second inequality follows from (4), (**) and (5), the third inequality follows from (2) and an iterated application of (3), and the last one from (*).

The following Borel-Cantelli Lemma is well-known (see for instance [Spr]).

Theorem 3.2 *Let (Z, ν) be a probability space. Let $(A_n)_{n\in\mathbb{N}}$ be a sequence of measurable subsets of Z such that there exists a constant $c > 0$ with $\nu(A_n \cap A_m) \leq c\nu(A_n)\nu(A_m)$ for every distinct integers n, m. Let $A_\infty = \bigcap_{n\in\mathbb{N}} \bigcup_{k\geq n} A_k$. Then $\nu(A_\infty) > 0$ if and only if $\displaystyle\sum_{n=0}^{\infty} \nu(A_n)$ diverges.* □

The result then follows. □

4 Spiraling geodesics

Let X be a proper CAT(-1) geodesic metric space. Let Γ be a non elementary discrete group of isometries of X, with finite critical exponent δ. Let Γ_0 be an almost malnormal convex-cocompact subgroup of infinite index in Γ with critical exponent δ_0, and let $C_0 = \mathcal{C}\Gamma_0$. Let $\pi_{C_0} : X \cup \partial_\infty X \to C_0 \cup \partial_\infty C_0$ be the closest point map. By Lemma 2.5, the number δ_0 belongs to $[0, \delta[$. Moreover, it follows from Section 2 that C_0 is non compact and that Γ_0 is the stabilizer in Γ of C_0.

Examples.

(1) Let γ_0 be an hyperbolic element of Γ, let C_0 be its translation axis and let Γ_0 be the stabilizer of C_0 (which is virtually infinite cyclic, and infinite cyclic when Γ is torsion free). Since Γ is non elementary, the subgroup Γ_0 has infinite index. Furthermore, if $\gamma \in \Gamma$ and $\gamma\Gamma_0\gamma^{-1} \cap \Gamma_0$ is infinite, then γ conjugates some hyperbolic element of Γ_0 to another one. The image by an element γ in Γ of the translation axis of an hyperbolic element α of Γ is the translation axis of $\gamma\alpha\gamma^{-1}$. Hence γ preserves C_0, therefore belongs to Γ_0. Therefore Γ_0 is an almost malnormal convex-cocompact subgroup of infinite index in Γ with critical exponent $\delta_0 = 0$.

(2) Let M be a complete Riemannian manifold with dimension $n \geq 2$ and sectional curvature at most -1, and $\pi : X \to M$ be a universal Riemannian covering, with covering group Γ. Let M_0 be a compact connected embedded totally geodesic submanifold in M of dimension k with $1 \leq k \leq n-1$, let C_0 be a connected component of the premimage of M_0 in X, and let Γ_0 be the stabilizer of C_0 in Γ (with good choices of base points, Γ can be identified with the fundamental group of M, and Γ_0 with the image in the fundamental

group of M of the fundamental group of M_0). Then Γ_0 is an almost malnormal (for instance by Proposition 2.6 (3)) convex-cocompact subgroup of infinite index in Γ. If M has constant sectional curvature -1, then $\delta = n-1$ and $\delta_0 = k-1$.

(3) Let $X = \mathbb{H}^3_{\mathbb{R}}$ be the real hyperbolic space of dimension 3, and Γ be a Kleinian group. If Γ_0 is a precisely invariant quasi-fuschian subgroup, without parabolic elements, of infinite index in Γ, then Γ_0 is an almost malnormal (by Proposition 2.6 (2)) convex-cocompact subgroup of infinite index in Γ.

After these examples, let us proceed. Denote by R_0 the set of double cosets

$$R_0 = \Gamma_0 \backslash (\Gamma - \Gamma_0) / \Gamma_0 \;.$$

For every $r = [\gamma]$ in R_0, define

$$D(r) = d(C_0, \gamma C_0) \in [0, +\infty[\;,$$

which does not depend on the representative γ of r. The next result says that the subset $\{D(r) \;:\; r \in R_0\}$ of $[0, +\infty[$ is discrete, with finite multiplicities.

Lemma 4.1 *For every $c \geq 0$, the set of elements r in R_0 such that $D(r) \leq c$ is finite.*

Proof. For every $c \geq 0$, assume that there exists a sequence of pairwise distinct elements $([\gamma_i])_{i\in\mathbb{N}}$ in R_0 such that $D([\gamma_i]) \leq c$ for every i. Fix x_* in C_0, and let D be the diameter of $\Gamma_0\backslash C_0$. For every i in $\mathbb{N}$, let x_i in C_0 and y_i in $\gamma_i C_0$ be any points such that $d(x_i, y_i) \leq c+1$. Up to replacing γ_i by another representative of $[\gamma_i]$, we may assume that $d(x_i, x_*) \leq D$ and $d(y_i, \gamma_i x_*) \leq D$. Hence $d(x_*, \gamma_i x_*) \leq 2D + c + 1$ for every i, which contradicts the discreteness of Γ. □

Proposition 4.2 *Assume that* Card $\Gamma x \cap B(x, n) \asymp e^{\delta n}$ *for some (hence every) x in X. Then there exists N in $\mathbb{N} - \{0\}$ such that*

$$\mathrm{Card}\;\{r \in R_0 \;:\; n \leq D(r) < n + N\} \asymp e^{\delta n} \;.$$

Proof. As $\delta_0 < \delta$, the proof is the same as the proof of [HP4, Theo. 3.4], up to replacing the horoball HB_0 by C_0. □

Define $X_0 = \Gamma_0 \backslash X$, and $\partial_\infty X_0 = \Gamma_0 \backslash (\partial_\infty X - \Lambda\Gamma_0)$. Since $\Gamma_0\backslash C_0$ is compact, and since the closest point map is a continuous Γ_0-equivariant map from $\partial_\infty X - \Lambda\Gamma_0$ to C_0, the space $\partial_\infty X_0$ is compact. The distance-like map d_{C_0} on $\partial_\infty X - \Lambda\Gamma_0$ is invariant under Γ_0, and we denote by d_0 the quotient distance-like map on $\partial_\infty X_0$, i.e.

$$d_0(\overline{x}, \overline{y}) = \inf_{x\in\overline{x},\; y\in\overline{y}} d_{C_0}(x, y) \;.$$

Let $r = [\gamma]$ be an element in R_0. Define Λ_r (which does not depend on the representative γ of r) as the image of $\gamma\Lambda\Gamma_0$ by the canonical projection $\partial_\infty X - \Lambda\Gamma_0 \rightarrow \partial_\infty X_0$. By Proposition 2.6 (2), it follows that $(\Lambda_r)_{r\in R_0}$ is a family of pairwise disjoint compact subsets of $\partial_\infty X_0$. For every $\epsilon > 0$, define $\mathcal{N}_r(\epsilon)$ as the ϵ-neigbourhood of Λ_r in $\partial_\infty X_0$ for the distance-like map d_0. Note that $\mathcal{N}_r(\epsilon) \subset \mathcal{N}_r(\epsilon')$ if $\epsilon < \epsilon'$.

Let $(\mu_x)_{x\in X}$ be a conformal density of dimension δ for Γ. Fix a base point x_0 in C_0. Define

$$\widetilde{\mu}_{\Gamma_0 x_0} = \sum_{\alpha\in\Gamma_0} \mu_{\alpha x_0}\ .$$

Lemma 4.3 *The map $\widetilde{\mu}_{\Gamma_0 x_0}$ is a locally finite Borel measure on $\partial_\infty X - \Lambda\Gamma_0$, which is invariant under Γ_0, and absolutely continuous with respect to the restriction to $\partial_\infty X - \Lambda\Gamma_0$ of μ_x for every x in X.*

We denote by $\mu_{\Gamma_0 x_0}$ the finite Borel measure on the compact quotient $\partial_\infty X_0$ of $\partial_\infty X - \Lambda\Gamma_0$ defined by $\widetilde{\mu}_{\Gamma_0 x_0}$.

Proof. Denote by $s\mapsto P_{x_0,\Gamma_0}(s) = \sum_{\alpha\in\Gamma_0}\ e^{-sd(\alpha x_0, x_0)}$ the Poincaré series of Γ_0 with base point x_0.

Let ξ be in $\partial_\infty X - \Lambda\Gamma_0$ and α be in Γ_0. The point αx_0 belongs to C_0. Hence the horosphere centered at ξ passing through $\pi_{C_0}(\xi)$ meets the geodesic ray from αx_0 to ξ in a point u. As C_0 is convex and $\pi_{C_0}(\xi)$ is the closed point in C_0 to ξ, by an easy CAT(-1) comparison argument, the distance $d(u,\pi_{C_0}(\xi))$ is at most 1 (and even $\log\frac{3+\sqrt{5}}{2}$). By the triangle inequality,



$$\beta_\xi(\alpha x_0, \pi_{C_0}(\xi)) = d(\alpha x_0, u) \geq d(\alpha x_0, \pi_{C_0}(\xi)) - d(u,\pi_{C_0}(\xi))\ .$$

Therefore

$$\begin{aligned}\beta_\xi(\alpha x_0, x_0) = \beta_\xi(\alpha x_0,\pi_{C_0}(\xi)) - \beta_\xi(x_0,\pi_{C_0}(\xi)) &\geq d(\alpha x_0,\pi_{C_0}(\xi)) - 1 - d(x_0,\pi_{C_0}(\xi))\\ &\geq d(\alpha x_0, x_0) - 1 - 2d(x_0,\pi_{C_0}(\xi))\ ,\end{aligned}$$

where the last inequation is again obtained by the triangle inequality. Hence

$$\sum_{\alpha\in\Gamma_0} \frac{d\mu_{\alpha x_0}}{d\mu_{x_0}}(\xi) = \sum_{\alpha\in\Gamma_0}\ e^{-\delta\beta_\xi(\alpha x_0, x_0)} \leq e^{1+2d(x_0,\pi_{C_0}(\xi))}\ P_{x_0,\Gamma_0}(\delta)\ .$$

The right hand side, as $\delta > \delta_0$, is a positive continous map of $\xi\in\partial_\infty X - \Lambda\Gamma_0$. Hence $\widetilde{\mu}_{\Gamma_0 x_0}$ is a locally finite Borel measure on $\partial_\infty X - \Lambda\Gamma_0$. It is clearly invariant under Γ_0 by construction and the equivariance property of $(\mu_x)_{x\in X}$. As $\beta_\xi(\alpha x_0, x_0) \leq d(\alpha x_0, x_0)$, we have, for every ξ in $\partial_\infty X - \Lambda\Gamma_0$,

$$P_{x_0,\Gamma_0}(\delta) \leq \frac{d\widetilde{\mu}_{\Gamma_0 x_0}}{d\mu_{x_0}}(\xi) \leq e^{1+2d(x_0,\pi_{C_0}(\xi))}\ P_{x_0,\Gamma_0}(\delta)\ , \tag{- 3 -}$$

hence $\widetilde{\mu}_{\Gamma_0 x_0}$ and μ_{x_0} have the same measure class on $\partial_\infty X - \Lambda\Gamma_0$. □

Theorem 4.4 *There exist constants $c, c' > 0$ such that, for every r in R_0 and ϵ in $]\,0, c' e^{-D(r)}]$,*

$$\frac{1}{c}\ e^{-\delta_0 D(r)}\epsilon^{\delta-\delta_0} \leq \mu_{\Gamma_0 x_0}(\mathcal{N}_r(\epsilon)) \leq c\ e^{-\delta_0 D(r)}\epsilon^{\delta-\delta_0}\ .$$

Proof. For every double coset r in R_0, choose a representative γ_r of r such that

$$d(x_0, \gamma_r x_0) = \min_{\alpha, \alpha' \in \Gamma_0} d(x_0, \alpha\gamma_r\alpha' x_0)\ .$$

Denote by $\mathcal{N}_{\epsilon', d'}(A)$ the (closed) ϵ'-neighbourhood of a subset A for a distance or a distance-like map d'. The subset $\mathcal{N}_{\epsilon, d_{C_0}}(\gamma_r\Lambda\Gamma_0)$ of $\partial_\infty X$ is compact. By Lemma 2.3, it is contained in $\partial_\infty X - \partial_\infty C_0$, and its diameter for the distance-like map d_{C_0} tends to 0 as $D(r)$ tends to $+\infty$ and ϵ tends to 0. Recall that Γ_0 acts isometrically and properly on $\partial_\infty X - \partial_\infty C_0$ for the distance-like map d_{C_0}. Hence there exists $N' \in \mathbb{N}$ and $c'_1 > 0$ such that for every ϵ in $]0, c'_1]$, for every r in R_0, we have

$$\mathrm{Card}\{\alpha \in \Gamma_0\ :\ \alpha\, \mathcal{N}_{\epsilon, d_{C_0}}(\gamma_r\Lambda\Gamma_0) \cap \mathcal{N}_{\epsilon, d_{C_0}}(\gamma_r\Lambda\Gamma_0) \neq \emptyset\} \leq N'\ .$$

By the construction of $\mu_{\Gamma_0 x_0}$, we have, for every r in R_0 and ϵ in $]0, c'_1]$,

$$\frac{1}{N'}\, \widetilde{\mu}_{\Gamma_0 x_0}(\mathcal{N}_{\epsilon, d_{C_0}}(\gamma_r\Lambda\Gamma_0)) \leq \mu_{\Gamma_0 x_0}(\mathcal{N}_r(\epsilon)) \leq \widetilde{\mu}_{\Gamma_0 x_0}(\mathcal{N}_{\epsilon, d_{C_0}}(\gamma_r\Lambda\Gamma_0))\ . \tag{- 4 -}$$

As $\Gamma_0\backslash C_0$ is compact and by the definition of the representatives γ_r, there exists $c'_2 > 0$ such that, for every r in R_0, for every $x \in \gamma_r(C_0 \cup \partial_\infty C_0)$, the closest point to x on C_0 is at distance at most c'_2 from x_0 (see also [HP4, Lem. 3.5]).

Hence, there exists a compact subset K of $\partial_\infty X - \partial_\infty C_0$ which contains $\gamma_r\Lambda\Gamma_0$ for every r in R_0. By Lemma 2.3 (2), there exists a compact subset K' of $\partial_\infty X - \partial_\infty C_0$ which contains $\mathcal{N}_{\epsilon, d_{C_0}}(\gamma_r\Lambda\Gamma_0)$ for every ϵ in $]0, c'_1]$ and every r in R_0. Hence by Lemma 2.3 (1), there exist two constants $c_3^\pm > 0$ such that for every r in R_0 and $\epsilon \in\,]0, c'_1]$,

$$\mathcal{N}_{c_3^- \epsilon, d_{x_0}}(\gamma_r\Lambda\Gamma_0) \subset \mathcal{N}_{\epsilon, d_{C_0}}(\gamma_r\Lambda\Gamma_0) \subset \mathcal{N}_{c_3^+ \epsilon, d_{x_0}}(\gamma_r\Lambda\Gamma_0)\ . \tag{- 5 -}$$

As K' and $\partial_\infty C_0$ are compact and disjoint, if c'_1 is small enough, then there exists a compact subset K'' of $\partial_\infty X - \partial_\infty C_0$ containing $\mathcal{N}_{c_3^+ \epsilon, d_{x_0}}(\gamma_r\Lambda\Gamma_0)$ (and hence $\mathcal{N}_{c_3^- \epsilon, d_{x_0}}(\gamma_r\Lambda\Gamma_0)$) for every r in R_0 and ϵ in $]0, c'_1]$. By the continuity of π_{C_0}, there exists a constant $c'_4 > 0$ such that for every r in R_0 and ϵ in $]0, c'_1]$, the subset $\pi_{C_0}(\mathcal{N}_{c_3^\pm \epsilon, d_{C_0}}(\gamma_r\Lambda\Gamma_0))$ is contained in the ball of center x_0 and radius c'_4.

By the definition of the representatives γ_r, for every r in R_0, for every $\xi \in \gamma_r\partial_\infty C_0$, the point $\gamma_r x_0$ is at distance at most a constant from the geodesic between x_0 and ξ (see also [HP4, Lem. 3.5]). Recall that for every η, η' in $\partial_\infty X$, if $d_{x_0}(\eta, \eta') \leq \epsilon'$, then the geodesic rays $[x_0, \eta[$ and $[x_0, \eta'[$ remain at distance bounded by a universal constant at least during a time $-\log\epsilon'$. Hence, if $c' \leq c'_1$ is small enough and $\epsilon \leq c' e^{-D(r)}$, then every geodesic ray from x_0 to a point ξ in $\mathcal{N}_{c_3^\pm \epsilon, d_{x_0}}(\gamma_r\Lambda\Gamma_0)$ passes at distance less than a constant from $\gamma_r x_0$. This has two consequences.

• First, using the change of base point formula for the visual distances, there exist two constants $c_5^\pm > 0$ such that for every $\epsilon \leq c'\, e^{-D(r)}$,

$$\mathcal{N}_{c_3^+ \epsilon, d_{x_0}}(\gamma_r\Lambda\Gamma_0) \subset \mathcal{N}_{c_5^+ e^{D(r)}\epsilon, d_{\gamma_r x_0}}(\gamma_r\Lambda\Gamma_0) \ \text{ and }\ \mathcal{N}_{c_5^- e^{D(r)}\epsilon, d_{\gamma_r x_0}}(\gamma_r\Lambda\Gamma_0) \subset \mathcal{N}_{c_3^- \epsilon, d_{x_0}}(\gamma_r\Lambda\Gamma_0)\ . \tag{- 6 -}$$

• Second, for every ξ in $\mathcal{N}_{c_3^{\pm}\epsilon,d_{x_0}}(\gamma_r\Lambda\Gamma_0)$, the number $|\beta_\xi(x_0,\gamma_r x_0)-d(x_0,\gamma_r x_0)|$ is bounded by a constant. Hence there exist constants $c_6^{\pm}>0$ such that for every r in R_0 and for every ξ in $\mathcal{N}_{c_3^{\pm}\epsilon,d_{x_0}}(\gamma_r\Lambda\Gamma_0)$,

$$c_6^-\ e^{-\delta D(r)} \leq \frac{d\mu_{x_0}}{d\mu_{\gamma_r x_0}}(\xi) \leq c_6^+\ e^{-\delta D(r)}\ . \tag{-7-}$$

By the Radon-Nykodim derivative estimates in Equation (-3-) and the definition of c_4', there exist constants $c_7^{\pm}>0$ such that for every ϵ in $]0,c_1']$, every r in R_0, and every ξ in $\mathcal{N}_{c_3^{\pm}\epsilon,d_{x_0}}(\gamma_r\Lambda\Gamma_0)$,

$$c_7^- \leq \frac{d\widetilde{\mu}_{\Gamma_0 x_0}}{d\mu_{x_0}}(\xi) \leq c_7^+\ . \tag{-8-}$$

By Sullivan's shadow lemma (see for instance [Rob, Lem. 1.3]), for every constant $c_8'>0$ big enough, there exist constants $c_9^{\pm}>0$ such that, for every γ in Γ,

$$c_9^-e^{-\delta d(x_0,\gamma x_0)} \leq \mu_{x_0}({}_{x_0}B(\gamma x_0,c_8')) \leq c_9^+e^{-\delta d(x_0,\gamma x_0)}\ . \tag{-9-}$$

For every $t\geq 0$, define $\Gamma_0[t]=\{\alpha\in\Gamma_0\ :\ d(x_0,\alpha x_0)\leq t\}$. For every $\epsilon'\in\,]0,1]$ and $\kappa>0$, define

$$A^+_{\epsilon',\kappa}=\Gamma_0[-\log\epsilon'+\kappa]-\Gamma_0[-\log\epsilon'-\kappa]\ \text{ and }\ A^-_{\epsilon',\kappa}=\Gamma_0[-\log\epsilon'+2\kappa]-\Gamma_0[-\log\epsilon'+\kappa]\ .$$

Let $\epsilon'\in\,]0,1]$, $\eta\in\partial_\infty X$ and $\eta'\in\Lambda\Gamma_0$ be such that $d_{x_0}(\eta,\eta')\leq\epsilon'$. Let u be the point of $[x_0,\eta[$ at distance $-\log\epsilon'$ from x_0. By the definition of d_{x_0} and the properties of the geodesic rays in a CAT(-1) metric space, there exists a universal constant c_8'' such that η' belongs to ${}_{x_0}B(u,c_8'')$. Let $c_{10}^+>0$ be at least the (finite) diameter of $\Gamma_0\backslash C_0$. Since $\partial_\infty C_0=\Lambda\Gamma_0$ and by convexity, the geodesic ray $[x_0,\eta[$ is contained in C_0. Hence there exists α in Γ_0 such that $d(u,\alpha x_0)\leq c_{10}^+$. Let c_8' be big enough (at least $c_8''+c_{10}^+$ and such that Equation (-9-) holds). Then, by the triangle inequality, $B(u,c_8'')$ is contained in $B(\alpha x_0,c_8')$. Note that $-\log\epsilon'-c_{10}^+\leq d(x_0,\alpha x_0)\leq-\log\epsilon'+c_{10}^+$. Therefore, for every ϵ' in $]0,1]$, we have

$$\mathcal{N}_{\epsilon',d_{x_0}}(\Lambda\Gamma_0)\subset\bigcup_{\alpha\in A^+_{\epsilon',c_{10}^+}}{}_{x_0}B(\alpha x_0,c_8')\ . \tag{-10-}$$

As Γ_0 is convex-cocompact, there exists a constant $c_8'''>0$ such that for every α in Γ_0, the segment $[x_0,\alpha x_0]$ is at distance at most c_8''' from a geodesic ray starting from x_0 and contained in C_0. Assume that $c_{10}^->0$ is at least $c_8'''+c_8'$. Let $\epsilon'\in\,]0,1]$, $\alpha\in A^-_{\epsilon',c_{10}^-}$ and $\eta'\in{}_{x_0}B(\alpha x_0,c_8')$. Let v be a point on $[x_0,\eta'[$ at distance at most c_8' from αx_0. Let $\eta\in\partial_\infty C_0$ and $u\in[x_0,\eta[$ be such that $d(u,\alpha x_0)\leq c_8'''$, which exist by the definition of c_8'''. Then by the definition of d_{x_0} and the triangle inequality, we have

$$d_{x_0}(\eta,\eta')\leq e^{\frac{1}{2}(d(u,v)-d(x_0,u)-d(x_0,v))}\leq e^{c_8'+c_8'''-d(x_0,\alpha x_0)}\leq\epsilon'\ ,$$

since $\alpha\in A^-_{\epsilon',c_{10}^-}$. Therefore, for every ϵ' in $]0,1]$, we have

$$\bigcup_{\alpha\in A^-_{\epsilon',c_{10}^-}}{}_{x_0}B(\alpha x_0,c_8')\subset\mathcal{N}_{\epsilon',d_{x_0}}(\Lambda\Gamma_0)\ .$$

If c^+_{10} and then c^-_{10} are big enough, as Γ_0 is convex-cocompact (by for instance [Rob] if Γ_0 is non elementary, and even if $\delta_0 = 0$, since then, by the assumptions, Γ_0 contains an hyperbolic element generating a finite index (infinite cyclic) subgroup), note that there exist constants $c^\pm_{11} > 0$ such that for every ϵ' in $]0,1]$, we have

$$\mathrm{Card}\; A^-_{\epsilon',c^-_{10}} \geq c^-_{11}\,(\epsilon')^{-\delta_0} \;\text{ and }\; \mathrm{Card}\; A^+_{\epsilon',c^+_{10}} \leq c^+_{11}\,(\epsilon')^{-\delta_0}\;. \tag{- 11 -}$$

Let $A^*_{\epsilon'}$ be a maximal subset of $A^-_{\epsilon',c^-_{10}}$ such that the shadows ${}_{x_0}B(\alpha x_0, c'_8)$ for α in $A^*_{\epsilon'}$ are pairwise disjoint. By maximality, for every α in $A^-_{\epsilon',c^-_{10}}$, there exists α' in $A^*_{\epsilon'}$ such that αx_0 and $\alpha' x_0$ are at bounded distance. Hence there exists a constant $c'_{12} > 0$ such that $\mathrm{Card}\; A^*_{\epsilon'} \geq c'_{12}\;\mathrm{Card}\; A^-_{\epsilon',c^-_{10}}$.

Let us now prove the upper bound in Theorem 4.4. Let r in R_0 and ϵ in $]\,0, c'\,e^{-D(r)}]$. Note that if $c' > 0$ is small enough, then $\epsilon \leq c'_1$ and $c^\pm_5\,e^{D(r)}\epsilon \leq c^\pm_5\,c' \leq 1$. We have

$$\begin{aligned}
\mu_{\Gamma_0 x_0}(\mathcal{N}_r(\epsilon)) &\leq \widetilde{\mu}_{\Gamma_0 x_0}(\mathcal{N}_{\epsilon,d_{C_0}}(\gamma_r\Lambda\Gamma_0)) \leq \widetilde{\mu}_{\Gamma_0 x_0}(\mathcal{N}_{c^+_3\epsilon,d_{x_0}}(\gamma_r\Lambda\Gamma_0)) && \text{by (- 4 -) and (- 5 -)}\\
&\leq c^+_7\,\mu_{x_0}(\mathcal{N}_{c^+_3\epsilon,d_{x_0}}(\gamma_r\Lambda\Gamma_0)) && \text{by (- 8 -)}\\
&\leq c^+_7\,c^+_6\,e^{-\delta D(r)}\mu_{\gamma_r x_0}(\mathcal{N}_{c^+_5 e^{D(r)}\epsilon,d_{\gamma_r x_0}}(\gamma_r\Lambda\Gamma_0)) && \text{by (- 7 -) and (- 6 -)}\\
&= c^+_7\,c^+_6\,e^{-\delta D(r)}\mu_{x_0}(\mathcal{N}_{c^+_5 e^{D(r)}\epsilon,d_{x_0}}(\Lambda\Gamma_0)) && \text{by invariance}\\
&\leq c^+_7\,c^+_6\,e^{-\delta D(r)}\sum_{\alpha\in A^+_{c^+_5 e^{D(r)}\epsilon,\,c^+_{10}}}\mu_{x_0}({}_{x_0}B(\alpha x_0, c'_8)) && \text{by (- 10 -)}\\
&\leq c^+_7\,c^+_6\,e^{-\delta D(r)}c^+_9\,e^{-\delta(-\log(c^+_5 e^{D(r)}\epsilon)-c^+_{10})}c^+_{11}(c^+_5\,e^{D(r)}\epsilon)^{-\delta_0} && \text{by (- 9 -) and (- 11 -)}\\
&= c^+_{13}\,e^{-\delta_0 D(r)}\epsilon^{\delta-\delta_0}\;,
\end{aligned}$$

for some constant $c^+_{13} > 0$, which proves the upper bound.

Similarly for the lower bound,

$$\begin{aligned}
\mu_{\Gamma_0 x_0}(\mathcal{N}_r(\epsilon)) &\geq \frac{1}{N'}\;\widetilde{\mu}_{\Gamma_0 x_0}(\mathcal{N}_{\epsilon,d_{C_0}}(\gamma_r\Lambda\Gamma_0)) \geq \frac{1}{N'}\;\widetilde{\mu}_{\Gamma_0 x_0}(\mathcal{N}_{c^-_3\epsilon,d_{x_0}}(\gamma_r\Lambda\Gamma_0))\\
&\geq \frac{c^-_7\,c^-_6}{N'}\;e^{-\delta D(r)}\mu_{x_0}(\mathcal{N}_{c^-_5 e^{D(r)}\epsilon,d_{x_0}}(\Lambda\Gamma_0))\\
&\geq \frac{c^-_7\,c^-_6}{N'}\;e^{-\delta D(r)}\mu_{x_0}\Big(\bigcup_{\alpha\in A^*_{c^-_5 e^{D(r)}\epsilon}}{}_{x_0}B(\alpha x_0, c'_8)\Big)\\
&= \frac{c^-_7\,c^-_6}{N'}\;e^{-\delta D(r)}\sum_{\alpha\in A^*_{c^-_5 e^{D(r)}\epsilon}}\mu_{x_0}({}_{x_0}B(\alpha x_0, c'_8))\\
&\geq \frac{c^-_7\,c^-_6}{N'}\;e^{-\delta D(r)}\;c^-_9\;e^{-\delta(-\log(c^-_5 e^{D(r)}\epsilon)+2c^-_{10})}\;c'_{12}\;c^-_{11}(c^-_5\,e^{D(r)}\epsilon)^{-\delta_0}\\
&= c^-_{13}\,e^{-\delta_0 D(r)}\epsilon^{\delta-\delta_0}\;,
\end{aligned}$$

for some constant $c^-_{13} > 0$, which proves the result. □

Lemma 4.5 *For every $N \in \mathbb{N} - \{0\}$, there exists $c'' > 0$ such that for every n in $\mathbb{N}$, for every distinct r and r' in R_0 such that $D(r)$ and $D(r')$ belong to $[nN, (n+1)N[$, the subsets $\mathcal{N}_r(c''\, e^{-nN})$ and $\mathcal{N}_{r'}(c''\, e^{-nN})$ are disjoint.*

Proof. Let $N \in \mathbb{N} - \{0\}$, and $c'' \leq 1$ be small enough, to be determined during the proof. Assume by absurd that there exists n in $\mathbb{N}$, distinct r and r' in R_0 such that $D(r), D(r') \in [nN, (n+1)N[$, and that the subsets $\mathcal{N}_r(c''\, e^{-nN})$ and $\mathcal{N}_{r'}(c''\, e^{-nN})$ have non empty intersection. Then, there exist representatives γ, γ' of the double cosets r, r' and points ξ, ξ' in $\gamma\Lambda\Gamma_0, \gamma'\Lambda\Gamma_0$ respectively, and an element η in $\partial_\infty X - \partial_\infty C_0$ which is different from ξ, ξ', such that $d_{C_0}(\xi, \eta)$ and $d_{C_0}(\xi', \eta)$ are at most $c''\, e^{-nN}$, and in particular at most c".

Since there are only finitely many r's with $D(r)$ less than a constant, and since the subsets $\gamma\Lambda\Gamma_0$ for γ in $(\Gamma - \Gamma_0)/\Gamma_0$ are pairwise disjoint (by Proposition 2.6 (2)) closed subsets, we may assume that $D(r)$ and $D(r')$ are bigger than any given constant $c''_1 > 0$. In particular, $D(r)$ and $D(r')$ are positive.

By Lemma 2.3 (4), there exists a universal constant $c''_2 \geq 1$ such that if $d_{C_0}(\eta', \eta'') \leq 1/c''_2$, then the geodesic line between η' and η'' is disjoint from C_0, and the length of the common perpendicular segment between $]\eta', \eta''[$ and C_0 is at most $-\log d_{C_0}(\eta', \eta'') + c''_2$ and at least $-\log d_{C_0}(\eta', \eta'') - c''_2$. Assume that $c'' \leq 1/c''_2$.



Let $p_\xi, p_{\xi'}, p_\eta$ be the closest point on C_0 to ξ, ξ', η respectively. Let $[x, y]$ (resp. $[x', y']$; $[x_\xi, y_\xi]$; $[x_{\xi'}, y_{\xi'}]$) be the common perpendicular between C_0 and γC_0 (resp. C_0 and $\gamma' C_0$; C_0 and $]\xi, \eta[$; C_0 and $]\xi', \eta[$), with $x, x', x_\xi, x_{\xi'}$ in C_0. Let z, z' be the closest point to y, y' on $[p_\xi, \xi[, [p_{\xi'}, \xi'[$ respectively. Let v, u, v', u' be the closest points to $y_\xi, y_\xi, y_{\xi'}, y_{\xi'}$ on $[p_\xi, \xi[, [p_\eta, \eta[, [p_{\xi'}, \xi'[, [p_\eta, \eta[$ respectively (see the picture on the right). Let $z_\xi, v_\gamma, w_{\gamma'}$ be the closest point to z, v, u on $[x_\xi, y_\xi], \gamma C_0, \gamma' C_0$ respectively. We have $d(x, y) = D(r)$, $d(x', y') = D(r')$.

Assume that $c'' \leq e^{-c''_1 - c''_2}$, so that $d(x_\xi, y_\xi)$, $d(x_{\xi'}, y_{\xi'})$ are at least $-\log(c'') - c''_2 \geq c''_1$. By the convexity of C_0 and quasi-geodesic arguments, if c''_1 is bigger than some universal constant, then there exists a universal constant c''_3 such that the distances $d(z, y)$, $d(x, p_\xi)$, $d(z', y')$, $d(x', p_{\xi'})$, $d(v, y_\xi)$, $d(u, y_\xi)$, $d(v', y_{\xi'})$, $d(u', y_{\xi'})$, $d(p_\xi, x_\xi)$, $d(p_\eta, x_\xi)$, $d(p_\eta, x_{\xi'})$, $d(p_{\xi'}, x_{\xi'})$ are at most c''_3. Hence

$$\begin{aligned} d(z, v) &\geq d(z, y_\xi) - d(y_\xi, v) \geq d(z_\xi, y_\xi) - c''_3 \geq d(y_\xi, x_\xi) - d(x_\xi, z_\xi) - c''_3 \\ &\geq (-\log(c'' e^{-nN}) - c''_2) - (d(x, y) + 2c''_3) - c''_3 \\ &\geq -\log c'' + nN - c''_2 - N(n+1) - 3\, c''_3 = -\log c'' - N - c''_2 - 3c''_3\ . \end{aligned}$$

In particular, if c'' is small, then the points p_ξ, z, v are in this order on $[p_\xi, \xi[$. So that by convexity $d(y, v_\gamma) \geq d(z, v) - c''_3 \geq -\log c'' - N - c''_2 - 4c''_3$, which is big if c'' is small. Similarly, $p_{\xi'}, z', v'$ are in this order on $[p_{\xi'}, \xi'[$.

Up to permuting ξ, ξ', we may assume that p_η, u, u', η are in this order on the geodesic ray $[p_\eta, \eta[$. By convexity properties of the distance, $d(u, w_{\gamma'}) \leq 3c''_3$. Hence as above, there exists a constant $c''_4 > 0$ such that $d(y', w_{\gamma'}) \geq -\log c'' - c''_4$.

Consider the geodesic hexagon with vertices $x, y, v_\gamma, w_{\gamma'}, y', x'$. We have $d(x, x') \leq 6c''_3$, $d(v_\gamma, w_{\gamma'}) \leq 6c''_3$, and $[x, y]$ (resp. $[x', y']$) is the shortest segment between points of $[x, x']$ and $[y, v_\gamma]$ (resp. $[y', w_{\gamma'}]$). Furthermore $d(x, y), d(x', y')$ may be assumed to be bigger than any constant and differ by at most a constant (that is by N), and $d(y, v_\gamma), d(y', w_{\gamma'})$ are bigger than $-\log c''$ minus a constant. For instance by using techniques of approximation by trees (see for example [GH, page 33]) on the above hexagon, the geodesic segments $[y, v_\gamma]$ and $[y', w_{\gamma'}]$, contained respectively in γC_0 and $\gamma' C_0$, are arbitrarily long if c'' is small enough; moreover their first endpoints and last endpoints are at bounded distance. Let $\epsilon = 1$ and $\kappa(\epsilon)$ be given by Proposition 2.6 (4). Hence by hyperbolicity, the ϵ-neighborhoods of γC_0 and $\gamma' C_0$ meet in a segment of length that can be made bigger than $\kappa(\epsilon)$ if c'' is small enough. This is a contradiction to Proposition 2.6 (4). $\Box$

A map $\psi : [0, +\infty[\, \to \,]0, +\infty[$ is called *slowly varying* (see [Sul]) if it is measurable and if there exist constants $B > 0$ and $A \geq 1$ such that for every x, y in $\mathbb{R}_+$, if $|x - y| \leq B$, then $\psi(y) \leq A\,\psi(x)$. Recall (see for instance [HP4, Sec. 5]) that this implies that ψ is locally bounded, hence it is locally integrable; also, if $\log \psi$ is Lipschitz, then ψ is slowly varying; and for every $N \in \mathbb{N} - \{0\}$ and $\epsilon > 0$, the series $\sum_{n=0}^{\infty} \psi(Nn)^\epsilon$ converges if and only if the integral $\int_0^\infty \psi(t)^\epsilon dt$ converges.

The following statement is the main technical step towards our Khintchine-type theorem for the spiraling of geodesic rays in $\Gamma \backslash X$ around $\Gamma_0 \backslash C_0$. It gives a 0-1 measure result for the approximation of points in the limit set of Γ by points of the orbit under Γ of the limit set of Γ_0.

Theorem 4.6 *Let $X, \Gamma, \Gamma_0, \delta, \delta_0, (\mu_x)_{x \in X}$ be as above. Assume furthermore, for some (hence any) x in X, that μ_x is ergodic for the action of Γ, and that* Card $\Gamma x \cap B(x, n) \asymp e^{\delta n}$. *Let $f : [0, +\infty[\, \to \,]0, +\infty[$ be a slowly varying map.*

If $\int_1^{+\infty} f(t)^{\delta - \delta_0}\, dt$ converges (resp. diverges), then $\mu_{\Gamma_0 x_0}$-almost no (resp. every) point of $\partial_\infty X_0$ belongs to infinitely many $\mathcal{N}_r\big(f(D(r)) e^{-D(r)}\big)$ where $r \in R_0$.

Remark. In addition to the hypotheses on X in this theorem, assume in this remark that X is a Riemannian manifold with constant sectional curvature -1, that Γ is convex-cocompact and that Γ_0 is the stabilizer of a geodesic line. Then up to some rewriting, this result is already known, see for instance [DMPV] or the recent [BV]. But even in this particular case, our techniques are very different from the ones of [DMPV, BV].

Proof. By a similar reduction as in [HP4, Lem. 5.2], we may assume that $f \leq 1$. Define $g = -\log f : [0, +\infty[\, \to [0, +\infty[$.

We apply Theorem 3.1 with $Z = \partial_\infty X_0$, $\mu = \mu_{\Gamma_0 x_0}$, $I = R_0$, and, for every r in R_0, $n \in \mathbb{N}$ and $\epsilon > 0$, with $B_r(\epsilon) = \mathcal{N}_r(\epsilon)$, $I_n = \{r \in R_0 \; : \; Nn \leq D(r) < N(n+1)\}$ where N is as in Proposition 4.2, and $n_r = E[\frac{D(r)}{N}]$ where E denotes the integer part. Define, for every n in $\mathbb{N}$ and $\epsilon > 0$,

$$f_1(n) = e^{\delta n N}, \quad f_2(n) = c_2\, e^{-nN}, \quad f_3(n) = c_2\, e^{-(nN + g(nN))}, \quad f_4(n) = c_2^{\delta_0 - \delta}\, e^{-\delta_0 n N} \;,$$

$$f_5(\epsilon) = \epsilon^{\delta - \delta_0},$$

where c_2 is a small enough positive constant. In particular, we assume that c_2 is less than $c'e^{-N}$, where c' is the constant defined in Theorem 4.4, and less than the constant c'' defined in Lemma 4.5. Note that

$$f_1(n) f_4(n) f_5(f_3(n)) = e^{-(\delta-\delta_0)g(Nn)} = f(Nn)^{\delta-\delta_0} \,.$$

Hence, as f is slowly varying, the series $\sum_{n\in\mathbb{N}} f_1(n)f_4(n)f_5(f_3(n))$ converges if and only if the integral $\int_1^{+\infty} f^{\delta-\delta_0}$ converges.

Note that $B_r(\epsilon)$ is measurable and non-decreasing in ϵ, and that I_n is finite by Lemma 4.1. Assumption (1) of Theorem 3.1 is satisfied since g is non negative. The assumptions (2) and (3) are easily verified. Assumption (4) follows from Proposition 4.2. Assumption (5) follows from Theorem 4.4 and the first assumption on c_2. Assumption (6) is satisfied by Lemma 4.5 and the second assumption on c_2.

Let us check that Assumption (7) of Theorem 3.1 is also satisfied.

Let $r, r' \in R_0$ with $n = n_r < m = n_{r'}$ such that $\mathcal{N}_r(f_3(n))$ and $\mathcal{N}_{r'}(f_3(m))$ meet. Hence, there exists representatives γ, γ' of r, r', points ξ, ξ' in $\gamma\Lambda\Gamma_0, \gamma'\Lambda\Gamma_0$ respectively, and $\eta \in \partial_\infty X - \Lambda\Gamma_0$ such that $d_{C_0}(\xi, \eta) \leq f_3(n)$ and $d_{C_0}(\xi', \eta) \leq f_3(m)$. Let us prove that there exists a big enough constant $\lambda > 0$ such that $\mathcal{N}_{r'}(f_2(m))$ is contained in $\mathcal{N}_r(\lambda f_3(n))$.

Recall that there are only finitely many r's with $D(r)$ less than a constant. As d_{C_0} is bounded on $\partial_\infty X_0 \times \partial_\infty X_0$, the ϵ-neighbourhood for d_{C_0} of any non-empty set covers $\partial_\infty X_0$ if ϵ is big enough. Hence we may assume, if λ is big enough, that $D(r)$ and $D(r')$ are bigger than any given constant $c_6 > 0$. In particular, $D(r)$ and $D(r')$ are positive.

Let $p_\xi, p_{\xi'}, p_\eta, x, y, x', y', x_\xi, y_\xi, x_{\xi'}, y_{\xi'}, z, z', v, u, v', u', v_\gamma$ be as in the proof of Lemma 4.5 and its picture. Let $v_{\gamma'}$ be the closest point to v' on $\gamma' C_0$.

As in the proof of Lemma 4.5, if c_6 is bigger than a universal constant and if c_2 is small enough, then there exists a universal constant c_7 such that the following distances $d(z, y)$, $d(x, p_\xi,)$, $d(z', y')$, $d(x', p'_\xi)$, $d(v, y_\xi)$, $d(u, y_\xi)$, $d(v', y_{\xi'})$, $d(u', y_{\xi'})$, $d(p_\xi, x_\xi)$, $d(p_\eta, x_\xi)$, $d(p_\eta, x_{\xi'})$, $d(p_{\xi'}, x_{\xi'})$, $d(v, v_\gamma)$, $d(v', v_{\gamma'})$ are at most c_7. Furthermore, p_ξ, z, v, ξ are in this order on $[p_\xi, \xi[$ and $p_{\xi'}, z', v', \xi'$ are in this order on $[p_{\xi'}, \xi'[$, and $d(y, v_\gamma), d(y', v_{\gamma'})$ may be taken bigger than any given constant if c_2 is small enough.

Say that a point p is *above* q (resp. *below* q *by at most some constant* $h > 0$*)* with respect to C_0 if $d(p, C_0) \geq d(q, C_0)$ (resp. $d(q, C_0) \geq d(p, C_0) \geq d(q, C_0) - h$). As $m > n$, the point y' is above y or below y by at most some universal constant. If the point y' was below u by more than some big constant, then, if c_2 is small enough, some long subsegment of $[y', v_{\gamma'}]$ would have its endpoints at distance at most a few c_7's from the endpoints of some subsegment of $[y, v_\gamma]$, and as in the end of the proof of Lemma 4.5, this would contradict Proposition 2.6 (4). Therefore the point y' is either above, or below only by a some constant, the point u and hence y_ξ. So that for every $\lambda'' > 0$, there exists $\lambda' > 0$ such that the shadow (seen from p_ξ) of the ball of center y_ξ and radius $\lambda' > 0$ contains the shadow of the ball of center y' and radius $\lambda'' > 0$. Note that if λ'' is big enough, then the shadow of $B(y', \lambda'')$ contains $\mathcal{N}_{r'}(f_2(m))$, as seen in the proof of Lemma 4.5. But if λ is big enough, then $\mathcal{N}_r(\lambda f_3(n))$ contains the shadow of $B(y_\xi, \lambda')$. Hence Assumption (7) of Theorem 3.1 follows.

Let E_f be the set of points of $\partial_\infty X_0$ which belong to infinitely many

$$B_r(f_3(n_r)) = \mathcal{N}_r\Big(c_3\ e^{-NE[\frac{D(r)}{N}]}\ f\big(NE[\tfrac{D(r)}{N}]\big)\Big)$$

for r in R_0, and similarly let E'_f be the set of points of $\partial_\infty X_0$ which belong to infinitely many $\mathcal{N}_r\big(f(D(r))\ e^{-D(r)}\big)$. As f is slowly varying, there exists a constant $c_8 \geq 1$ such that $E_{\frac{1}{c_8}f} \subset E'_f \subset E_{c_8 f}$. Hence in order to prove Theorem 4.6, we only have to prove that if $\int_1^{+\infty} f^{\delta-\delta_0}$ converges (resp. diverges), then $\mu_{\Gamma_0 x_0}(E_f) = 0$ (resp. $\mu_{\Gamma_0 x_0}({}^c E_f) = 0$). If this integral converges, then the result follows from Part [A] of Theorem 3.1.

In the divergence case, Part [B] of Theorem 3.1 implies that $\mu_{\Gamma_0 x_0}(E_f) > 0$. Using the ergodicity of μ_x in a similar way to the end of the proof of Theorem 5.1 of [HP4], and the fact that $\mu_x(\Lambda\Gamma_0) = 0$ as $\delta_0 < \delta$, it follows that E_f has full measure. □

Let us now proceed towards our main result, Theorem 4.9.

Let ϵ be a positive real number and let $g : [0, +\infty[\ \to [0, +\infty[$ be a map such that $t \mapsto f(t) = e^{-g(t)}$ is slowly varying. A geodesic line ℓ in X will be called (ϵ, g)-*Liouville* with respect to (Γ, Γ_0) if there exist a sequence $(t_n)_{n\in\mathbb{N}}$ of positive times converging to $+\infty$ and a sequence $(\gamma_n)_{n\in\mathbb{N}}$ of elements of Γ such that $\ell(t)$ belongs to $\mathcal{N}_\epsilon(\gamma_n C_0)$ for every t in $[t_n, t_n + g(t_n)]$. The following remark implies that up to changing g by an additive constant (or equivalently up to changing f by a multiplicative constant), being (ϵ, g)-Liouville does not depend on ϵ, and depends only on the asymptotic class of ℓ.

Remark 4.7 (1) Note that if $\epsilon' \geq \epsilon$ and $g' \leq g$, then a geodesic line which is (ϵ, g)-Liouville is (ϵ', g')-Liouville.

(2) Note that by the hyperbolicity properties of X, for every ϵ' in $]\,0, \epsilon]$, there exists a constant $c(\epsilon, \epsilon') \geq 0$ such that for every convex subset C of X and every geodesic line ℓ in X, if the length h of the intersection of ℓ and $\mathcal{N}_\epsilon C$ is at least $c(\epsilon, \epsilon')$, then the length of the intersection of ℓ and $\mathcal{N}_{\epsilon'} C$ is at least $h - c(\epsilon, \epsilon')$ (see [PP] for precise estimates). In particular, if $g \geq c(\epsilon, \epsilon')$, then a geodesic line which is (ϵ, g)-Liouville is $(\epsilon', g - c(\epsilon, \epsilon'))$-Liouville.

(3) Recall that two geodesic lines ℓ, ℓ' in X are *asymptotic* if $d(\ell(t), \ell')$ (or equivalently $d(\ell'(t), \ell)$) is bounded (or equivalently tends to 0) as t tends to $+\infty$. Note that ℓ, ℓ' are asymptotic if and only if their points at infinity $\ell(+\infty), \ell'(+\infty)$ are equal.

By the strict convexity of ϵ-neighborhoods of convex subsets of X, if ℓ is an (ϵ, g)-Liouville geodesic line, and ℓ' is a geodesic line which is asymptotic to ℓ, then ℓ' is $(2\epsilon, g)$-Liouville, as well as $(\epsilon, g - \eta)$-Liouville for every constant $\eta > 0$ such that $g \geq \eta$.

Let $\pi_0 : X \cup (\partial_\infty X - \partial_\infty C_0) \to X_0 \cup \partial_\infty X_0$ be the canonical projection. The next lemma shows the relation between the (geometric) Liouville property of a geodesic line and the fact that its point at infinity belongs to a limsup subset considered in Theorem 4.6.

Lemma 4.8 *There exists $c''' > 0$ such that for every geodesic line ℓ in X such that $\ell(+\infty) \notin \bigcup_{\gamma\in\Gamma} \gamma\, \partial_\infty C_0$,* *(1) if ℓ is (ϵ, g)-Liouville, then the point $\pi_0(\ell(+\infty))$ belongs to infinitely many subsets $\mathcal{N}_r(c''' \, f(D(r)) \, e^{-D(r)})$ for r in R_0;*

(2) if $\pi_0(\ell(+\infty))$ belongs to infinitely many subsets $\mathcal{N}_r(\frac{1}{c'''} \, f(D(r)) \, e^{-D(r)})$ for r in R_0, then ℓ is (ϵ, g)-Liouville.

Proof. (1) Assume that ℓ is (ϵ, g)-Liouville. Up to replacing ϵ by 2ϵ and ℓ by an asymptotic line, as $\ell(+\infty) \notin \partial_\infty C_0$ and by Remark 4.7 (3), we may assume that $\ell(0)$ is the closest point in C_0 to $\ell(+\infty)$.

Let $(t_n)_{n \in \mathbb{N}}$ be a sequence of positive times, converging to $+\infty$. Let $(\gamma_n)_{n \in \mathbb{N}}$ in Γ be such that $\ell(t) \in \mathcal{N}_\epsilon(\gamma_n C_0)$ for every $t \in [t_n, t_n + g(t_n)]$. As Γ acts properly on X, and as its subgroup Γ_0 acts cocompactly on C_0, the family $(\gamma C_0)_{\gamma \in (\Gamma - \Gamma_0)/\Gamma_0}$ is locally finite. Hence $d(\gamma_n C_0, C_0)$ tends to $+\infty$ as $n \to +\infty$ (otherwise $\ell(+\infty)$ would belong to $\gamma \; \partial_\infty C_0$ for some $\gamma \in \Gamma$). In particular, up to extracting a subsequence, $\gamma_n \notin \Gamma_0$ and with $r_n = [\gamma_n] \in R_0$, the r_n's are pairwise distinct. Furthermore, we may assume that ℓ enters $\mathcal{N}_\epsilon(\gamma_n C_0)$ at the time t_n. Let $[p_n, q_n]$ be the shortest segment between C_0 and $\gamma_n C_0$, with $p_n \in C_0$, so that $D(r_n) = d(p_n, q_n)$. Let x_n (resp. y_n) be a point of $\gamma_n C_0$ such that $d(x_n, \ell(t_n)) \leq \epsilon$ (resp. $d\big(y_n, \ell(t_n + g(t_n))\big) \leq \epsilon$).



As we have already seen (in the proof of Lemma 4.5), there exists a constant $c_1''' > 0$ such that $[q_n, y_n]$ is contained in the c_1'''-neighborhood of a geodesic ray $[q_n, \xi_n[$ with $\xi_n \in \gamma_n \partial_\infty C_0$, and such that for n big enough, $d(p_n, \ell(0)) \leq c_1'''$. By hyperbolicity, the distance between p_n and the closest point of C_0 to ξ_n is at most a constant. By arguments similar to the ones in the proof of Lemma 4.5, it is easy to prove that there exists a constant $c_2''' \geq 0$ such that $-\log d_{C_0}(\xi_n, \ell(+\infty)) \geq t_n + g(t_n) - c_2'''$ and (using the fact that ℓ enters in $\mathcal{N}_\epsilon(\gamma_n C_0)$ at time t_n) that $|t_n - d(p_n, q_n)| \leq c_2'''$. As f is slowly varying, there exists a constant $c''' \geq 1$ such that $d_{C_0}(\xi_n, \ell(+\infty)) \leq c''' f(D(r_n)) \; e^{-D(r_n)}$. This proves the first assertion.

(2) Assume now that there exist a sequence $(r_n = [\gamma_n])_{n \in \mathbb{N}}$ of pairwise distinct elements in R_0 and $\xi_n \in \gamma_n \partial_\infty C_0$ such that $d_{C_0}(\xi_n, \ell(+\infty)) \leq \frac{1}{c'''} \; f(D(r_n)) \; e^{-D(r_n)}$ for every n, for some $c''' \geq 1$ big enough, to be determined later on. Let us prove that ℓ is (ϵ, g)-Liouville. Up to replacing ϵ by $\frac{\epsilon}{2}$ and ℓ by an asymptotic line, we may assume as above that $\ell(0)$ is the closest point in C_0 to $\ell(+\infty)$.

By Lemma 4.1, we have that $D(r_n) = d(C_0, \gamma_n C_0)$ tends to $+\infty$ as $n \to +\infty$ (hence is positive for n big enough). As above, by hyperbolicity, the closest point of C_0 to ξ_n is at distance at most a constant from the closest point of C_0 to $\gamma_n C_0$. By hyperbolicity and the definition of d_{C_0}, there exists a constant $c_3''' \geq 0$ such that between the times $t = D(r_n)$ and $t = D(r_n) + g(D(r_n)) + \log c''' - c_3'''$, the geodesic ray ℓ is at distance at most c_3''' from $\gamma_n C_0$. Hence, as in Remark 4.7 (3), there exists a constant $c_4''' \geq 0$ such that both $\ell(D(r_n) + c_4''')$ and $\ell(D(r_n) + g(D(r_n)) + \log c''' - c_3''' - c_4''')$ are at distance at most ϵ from $\gamma_n C_0$. Hence if c''' is big enough, by setting $t_n = D(r_n) + c_4'''$ and as f is slowly varying, the second assertion follows. $\square$

Theorem 4.9 *Let X be a proper* $\mathrm{CAT}(-1)$ *geodesic metric space. Let Γ be a non elementary discrete group of isometries of X, with finite critical exponent δ, of divergent type. Let $\widetilde{\mu}_{BM}$ be its Bowen-Margulis measure. Assume that* $\mathrm{Card}\,\Gamma x \cap B(x,n) \asymp e^{\delta n}$, *for some $x \in X$. Let $(\Gamma_i)_{i\in I}$ be a finite family of almost malnormal convex-cocompact subgroups of infinite index in Γ with critical exponents $(\delta_i)_{i\in I}$. Let $\delta_0^+ = \sup_{i\in I}\delta_i$ and $\delta_0^- = \inf_{i\in I}\delta_i$. Let $g : [0,+\infty[\ \to\ [0,+\infty[$ be a map such that $t \mapsto f(t) = e^{-g(t)}$ is slowly varying, and let $\epsilon > 0$.*

If $\int_1^{+\infty} f(t)^{\delta-\delta_0^+}\,dt$ diverges (resp. $\int_1^{+\infty} f(t)^{\delta-\delta_0^-}\,dt$ converges), then $\widetilde{\mu}_{BM}$-almost every (resp. no) element of $\mathcal{G}X$ is (ϵ, g)-Liouville with respect to (Γ, Γ_i) for every (resp. some) $i \in I$.

Remark. (1) The result still holds for a countable family $(\Gamma_i)_{i\in I}$, under the assumption that $\delta_0^+ < \delta$. There are examples of $\Gamma\backslash X$ with X and Γ as in the above theorem, such that the upper bound of the critical exponents of the infinite index subgroups in Γ is equal to (resp. is strictly less than) the critical exponent of Γ, as for instance the closed real hyperbolic 3-manifolds fibering over the circle (resp. the closed quaternionic hyperbolic manifolds, see for instance [Leu]).

(2) Assume in this remark that X is a Riemannian manifold, that Γ is cocompact and torsion free, and that Γ_0 is the stabilizer of a geodesic line. This correspond to the hypotheses of Theorem 1.1 (that appear above it). Then there might be a simpler proof using symbolic coding, as indicated to us by V. Kleptsyn, using the fact that the geodesic flow of $\Gamma\backslash X$ is then conjugated to a suspension of a Bernoulli shift. But this requires some serious amount of work, since some geometric features are difficult, to say the least, to translate by the coding. In our general situation, no such coding is possible anyway.

Proof. Note that the divergence (resp. convergence) of the integral in the statement is unchanged if one replaces f by a scalar multiple of it. Also recall that $\mu_{x_0}(\bigcup_{i\in I,\ \gamma\in\Gamma}\gamma\Lambda\Gamma_i) = 0$, since $\delta_0 < \delta$ (see Lemma 2.5).

When the index set I has only one element, the result follows from Theorem 4.6, by considering the conformal density $(\mu_x)_{x\in X}$ of dimension δ for Γ that is used in the construction (recalled in Section 2) of $\widetilde{\mu}_{\mathrm{BM}}$ (which is ergodic since Γ is of divergent type), and by the lemmas 4.8, 4.3, 2.4.

Using the fact that finite or countable unions of sets of measure 0 have measure 0, the result for general I follows. $\square$

Using the three examples at the beginning of Section 4, the theorems 1.1, 1.3, 1.4 in the introduction follow.

Similarly, to prove Proposition 1.5 of the introduction, we apply Theorem 4.9 to $X = T$ the tree in the statement of Proposition 1.5, with $I = \{0\}$ and Γ_0 the stabilizer in Γ of a geodesic line in T mapping to the cycle C in the statement of Proposition 1.5. We replace the map g in Theorem 4.9 by g/L for the map g in Proposition 1.5 (the map $t \mapsto \exp(-g(t)/L)$ is still slowly varying). When Γ is cocompact and torsion-free (anyone of these two assumptions may be non satisfied), then the symbolic dynamics argument alluded to above works easily, and gives an alternate proof. But no such coding is easy in general, even for lattices as simple as $\mathrm{PSL}_2(\mathbb{F}_q[X])$ in $\mathrm{PSL}_2\big(\mathbb{F}_q((X))\big)$, see for instance [BP].

Many other applications are possible, we will only give the next one.

We refer to [GP] (see also [Bou2, HaP]) for the definitions and basic properties of an hyperbolic building, which in particular, when locally finite, is a proper CAT(-1) geodesic metric space. For instance, for every integers $p \geq 5, q \geq 3$, let (W_p, S_p) be the hyperbolic Coxeter system generated by the reflections on the sides of a right angled regular real hyperbolic p-gon; Bourdon's building $I_{p,q}$ is (see for instance [Bou2]) the unique (up to isomorphism) hyperbolic building of dimension 2, modeled on (W_p, S_p), and whose links of vertices are bipartite graphs on $q+q$ vertices. It has a cocompact lattice $\Gamma_{p,q}$ with presentation

$$\langle s_1, \dots, s_p \mid \forall\, i \in \mathbb{Z}/p\mathbb{Z} \;\; s_i^q = 1, [s_i, s_{i+1}] = 1 \rangle \;,$$

where $s_1, \dots, s_p$ are generators of the pointwise stabilizers of the p panels of a fundamental chamber $\mathfrak{C}$ of $I_{p,q}$. If q is even, let Γ_0 be the subgroup (isomorphic to W_p) generated by the elements $s_i^{\frac{q}{2}}$ for $1 \leq i \leq p$, which is, by the simple transitivity of the action of $\Gamma_{p,q}$ on the set of chambers, the stabilizer of a (unique) apartment $A_{\mathfrak{C}}$ in $I_{p,q}$ containing $\mathfrak{C}$.

Remark 4.10 *If q is even, then the subgroup Γ_0 is almost malnormal in $\Gamma_{p,q}$.*

Proof. (F. Haglund) Let V be the union of the closed chambers of I_{pq} meeting $A_{\mathfrak{C}}$, which is invariant by Γ_0. By convexity (and arguments as in Poincaré's theorem about reflection groups), the subgroup H of Γ generated by the pointwise stabilizers of the edges contained in the boundary of V has V as a strict fundamental domain, and is normalized by Γ_0. The subgroup Γ' of Γ generated by H and Γ_0, which is isomorphic to their semi-direct product, has finite index in Γ, since Γ is discrete and Γ_0 acts transitively on the chambers of $A_{\mathfrak{C}}$. Let Γ'' be a finite index torsion free subgroup of Γ' (which exists for instance since $\Gamma_{p,q}$ is linear, see for example [Kap]).

Let us prove that the stabilizer $S = \Gamma'' \cap \Gamma_0$ of $A_{\mathfrak{C}}$ in Γ'' is malnormal in Γ'', which proves the result. Assume by absurd that there exists γ in $\Gamma'' - S$ and s in $S - \{e\}$ such that s and $\gamma s \gamma^{-1}$ preserve $A_{\mathfrak{C}}$. By construction, two distinct translates of $A_{\mathfrak{C}}$ by elements of Γ' are disjoint. Hence $\gamma^{-1} A_{\mathfrak{C}}$ and $A_{\mathfrak{C}}$ are disjoint, and both preserved by s. The (unique) shortest segment between $\gamma^{-1} A_{\mathfrak{C}}$ and $A_{\mathfrak{C}}$ is then fixed by s, which contradicts the fact that Γ'' is torsion free. □

Corollary 4.11 *Let X be a locally finite thick hyperbolic building modeled on an hyperbolic Coxeter system (W, S). Let Γ be a cocompact lattice in the automorphism group of X with Bowen-Margulis measure μ. Let A be an appartment in X whose stabilizer Γ_A in Γ acts cocompactly on A and is almost malnormal in Γ. Denote by $k \geq 1$ the dimension of A (hence of X), and by δ the Hausdorff dimension of $\partial_\infty X$ (for any visual distance). Let $f \leq 1$ be a slowly varying map, and $\epsilon > 0$.*

If $\int_1^{+\infty} f(t)^{\delta - k + 1}\, dt$ converges (resp. diverges), then for μ-almost no (resp. every) ℓ in $\mathcal{G}X$, there exist positive times $(t_n)_{n \in \mathbb{N}}$ converging to $+\infty$ such that $\ell(t)$ belongs to $\Gamma \mathcal{N}_\epsilon A$ for every t in $[t_n, t_n - \log f(t_n)]$.

Proof. The apartments in an hyperbolic building are convex (for the CAT(-1) metric), hence Γ_A is convex-cocompact with critical exponent $k-1$. As X is thick, Γ_A has infinite index in Γ. The result follows from Theorem 4.9 (with I a singleton). □

Let us go back to the general situation of Theorem 4.9. The following result is a logarithm law-type result for the spiraling of geodesic lines in $\Gamma\backslash X$ around $\Gamma_0\backslash C_0$. For every $\epsilon > 0$ fixed, define the *penetration map* $\mathfrak{p} = \mathfrak{p}_{\Gamma\mathcal{N}_\epsilon C_0} : \mathcal{G}X \times [0, +\infty[\;\rightarrow [0, +\infty[$ in $\Gamma\mathcal{N}_\epsilon C_0$ of the geodesic lines in X, in the following way. For $(\ell, t) \in \mathcal{G}X \times [0, +\infty[$, if $\ell(t)$ does not belong to $\Gamma\mathcal{N}_\epsilon C_0$, then let $\mathfrak{p}(\ell, t) = 0$. Otherwise, let $\mathfrak{p}(\ell, t)$ be the upper bound of the lengths of the intervals I in $\mathbb{R}$ containing t such that there exists γ in Γ with $\ell(I)$ contained in $\gamma\mathcal{N}_\epsilon C_0$.

Theorem 4.12 *Let X be a proper CAT(-1) geodesic metric space. Let Γ be a non elementary discrete group of isometries of X, with finite critical exponent δ, of divergent type, and let $\widetilde{\mu}_{BM}$ be the Bowen-Margulis measure of Γ. Assume that* Card $\Gamma x \cap B(x, n) \asymp e^{\delta n}$, *for some $x \in X$. Let Γ_0 be an almost malnormal convex-cocompact subgroup of infinite index in Γ with critical exponent δ_0.*

Then for every $\epsilon > 0$, for $\widetilde{\mu}_{BM}$-almost every ℓ in $\mathcal{G}X$, we have

$$\limsup_{t\rightarrow+\infty} \frac{\mathfrak{p}(\ell, t)}{\log t} = \frac{1}{\delta - \delta_0}\;.$$

Proof. For every γ in Γ such that a geodesic line ℓ enters the ϵ-neighborhood of γC_0, let $t_{\ell,\gamma}$ be the entering time of ℓ in this neighborhood.

We apply Theorem 4.9 with $g_\kappa : t \mapsto \kappa \log(1 + t)$, which is a Lipschitz map $\mathbb{R}^+ \rightarrow \mathbb{R}^+$. Note that the integral $\int_1^{+\infty} t^{-(\delta-\delta_0)\kappa}\, dt$ diverges if and only if $\kappa \leq \frac{1}{\delta-\delta_0}$. If $\kappa_n = \frac{1}{\delta-\delta_0} + \frac{1}{n}$ for $n \in \mathbb{N} - \{0\}$, then the convergence part of Theorem 4.9 implies that for $\widetilde{\mu}_{\rm BM}$-almost every ℓ in $\mathcal{G}X$, for every γ in Γ such that ℓ meets $\gamma\mathcal{N}_\epsilon C_0$ with $t_{\ell,\gamma}$ big enough, we have $\mathfrak{p}(\ell, t_{\ell,\gamma}) \leq g_{\kappa_n}(t_{\ell,\gamma})$. Hence

$$\limsup_{t\rightarrow+\infty} \frac{\mathfrak{p}(\ell, t)}{\log t} = \limsup \frac{\mathfrak{p}(\ell, t_{\ell,\gamma})}{\log(1 + t_{\ell,\gamma})} \leq \kappa_n\;,$$

where the upper limit is taken on the $\gamma \in \Gamma - \Gamma_0$ such that ℓ meets $\gamma\mathcal{N}_\epsilon C_0$ and $t_{\ell,\gamma}$ tends to $+\infty$. As $n \rightarrow +\infty$, we get that $\limsup_{t\rightarrow+\infty} \frac{\mathfrak{p}(\ell,t)}{\log t} \leq \frac{1}{\delta-\delta_0}$. Similarly, using the divergence part of Theorem 4.9 with the function $g = g_\kappa$ where $\kappa = \frac{1}{\delta-\delta_0}$, we get that for $\widetilde{\mu}_{\rm BM}$-almost every ℓ in $\mathcal{G}X$, $\limsup_{t\rightarrow+\infty} \frac{\mathfrak{p}(\ell,t)}{\log t} \geq \frac{1}{\delta-\delta_0}$. $\square$

Corollary 1.2 in the introduction follows immediately.

5 Non-archimedean Diophantine approximation by quadratic irrational numbers

Let us now give an application of our results to Diophantine approximation in non-archimedian local fields.

Let $\widehat{K} = \mathbb{F}_q((X^{-1}))$ be the field of formal Laurent series in the variable X^{-1} over the finite field $\mathbb{F}_q$. Recall the definition of the absolute value of an element $f \in \widehat{K} - \{0\}$. Let $f = \sum_{i=n}^{\infty} a_i X^{-i}$ where $n \in \mathbb{Z}$ and $a_n \neq 0$. Then we define $\nu(f) = n$ and $|f|_\infty = q^{-\nu(f)}$. Endow the locally compact additive group $\widehat{K}$ with its (unique up to a constant factor) Haar measure μ. Let $K = \mathbb{F}_q(X)$.

Let $\mathbb{T}_q$ be the Bruhat-Tits tree of $(\mathrm{SL}_2, \widehat{K})$; we refer to [Ser] for any background on $\mathbb{T}_q$. Identify as usual $\partial_\infty \mathbb{T}_q$ and $\widehat{K} \cup \{\infty\}$, so that the action of $\mathrm{SL}_2(\widehat{K})$ on $\mathbb{T}_q$ extends continuously by the action by homographies of $\mathrm{SL}_2(\widehat{K})$ on $\widehat{K} \cup \{\infty\}$. Let x_0 be the standard base point in $\mathbb{T}_q$. Note that the Hausdorff dimension of the visual distance d_{x_0} is $\log q$, as $\mathbb{T}_q$ is a regular tree of degree $q+1$.

We refer for instance to [Las, Sch] for nice surveys of the Diophantine approximation properties of elements in $\widehat{K}$ by elements in K, a geometric interpretation of which being given in [Pau]. Here, we are interested in approximating elements of $\widehat{K}$ by elements in the set K_2 of irrational quadratic elements in $\widehat{K}$ over K. For every α in K_2, let α^* be its Galois conjugate (the other root of its minimal polynomial), and define its *height* by

$$h(\alpha) = |\alpha - \alpha^*|_\infty{}^{-1} \; .$$

We will not make precise here the relationship with the standard height of an element of the projective line over the algebraic closure of K, see for instance [HS].

Let $\Gamma = \mathrm{PSL}_2(\mathbb{F}_q[X]) = \mathrm{SL}_2(\mathbb{F}_q[X])/\{\pm \mathrm{id}\}$, which is a (non-uniform) lattice of $\mathbb{T}_q$ (see for instance [Ser]), hence a non-elementary discrete group of isometries of $\mathbb{T}_q$, whose critical exponent δ is equal to the Hausdorff dimension of d_{x_0}, that is $\delta = \log q$. See for instance [BP] for a (well known) proof that the restrictions to $\partial_\infty \mathbb{T}_q - \{\infty\} = \widehat{K}$ of the Patterson-Sullivan measures of Γ have the same measure class as the Haar measure μ of $\widehat{K}$.

Proof of Theorem 1.6. Let γ_0 be an hyperbolic element of Γ, C_0 be its translation axis in $\mathbb{T}_q$, and Γ_0 be the stabilizer of C_0 in Γ, which is convex-cocompact with critical exponent $\delta_0 = 0$. It is easy to verify that the set of points at infinity of C_0 is $\{\alpha, \alpha^*\}$ for some α in K_2; and that any such pair is the set of endpoints of some hyperbolic element of Γ (one can for instance use the fact that Artin's continued fraction expansion of an element in K_2 is eventually periodic (see for example [Las])).

Note that for every $\gamma \in \Gamma$ and $\alpha \in K_2$, the element $\gamma\alpha$ is still in K_2, $(\gamma\alpha)^* = \gamma\alpha^*$ and $\gamma\{\alpha, \alpha^*\} \cap \{\alpha, \alpha^*\} \neq \emptyset$ if and only if $\gamma \in \Gamma_0$.

Denote by d_∞ the Hamenstädt distance on $\partial_\infty \mathbb{T}_q - \{\infty\} = \widehat{K}$ defined by the horosphere centered at ∞ and passing through x_0. It is proved in [Pau, Coro. 5.2] that $d_\infty(\xi, \xi') = |\xi - \xi'|_\infty^{\frac{1}{\log q}}$, for every ξ, ξ' in $\widehat{K}$.

Lemma 5.1 *For every ξ_0 in $\widehat{K} - \{\alpha, \alpha_*\}$, there exists a neighborhood V of ξ_0 and a constant $c_* > 0$ such that for every ξ, ξ' in V,*

$$d_{C_0}(\xi, \xi') = c_*\, d_\infty(\xi, \xi') = c_*\, |\xi - \xi'|_\infty^{\frac{1}{\log q}} \; .$$

Furthermore, for every γ in $\Gamma - \Gamma_0$ such that $\gamma\alpha$ and $\gamma\alpha^*$ belongs to V, we have

$$e^{-D([\gamma])} = c_*\, |\gamma\alpha - \gamma\alpha^*|_\infty^{\frac{1}{\log q}} \; .$$

Proof. Let p_0 be the intersection of the geodesic line $]\infty, \xi_0[$ in $\mathbb{T}_q$ with the horosphere centered at ∞ passing through x_0. Let $q_0 = \pi_{C_0}(\xi_0)$, and $u_0 \in X$ such that $]\xi_0, \infty[\,\cap\,]\xi_0, q_0] =]\xi_0, u_0]$. In the picture on the right, we assume that C_0 and $]\infty, \xi_0[$ are disjoint, and that $p_0 \in [u_0, \infty[$. But the following reasoning is independent of these assumptions. Let $c_* = e^{-\beta_{\xi_0}(q_0,p_0)}$. If ξ, ξ' are close enough to ξ_0, and $]\xi, \infty[\,\cap\,]\xi', \infty[= [p, \infty[$, then $p_0, u_0 \in [p, \infty[$, $\pi_{C_0}(\xi) = \pi_{C_0}(\xi') = q_0$, $\beta_{\xi_0}(q_0, p_0) = d(q_0, p) - d(p_0, p)$, $d_\infty(\xi, \xi') = e^{-d(p_0,p)}$ and $d_{C_0}(\xi, \xi') = e^{-d(q_0,p)}$, hence the first result follows.

As $d_{C_0}(\gamma\alpha, \gamma\alpha^*) = e^{-D([\gamma])}$ if $\gamma\alpha, \gamma\alpha^*$ are closed enough to ξ_0 (see Equation (- 1 -)), the second result follows from the first one. □



Let $\varphi : [0, +\infty[\, \to\,]0, 1]$ be a map with $t \mapsto f(t) = \varphi(q^t)^{\frac{1}{\log q}}$ slowly varying, and let $g : t \mapsto -\log f(t) = -\log_q \varphi(q^t)$, so that $\varphi(t) = q^{-g(\log_q t)}$. By an easy change of variable, the integral $\int_1^{+\infty} \varphi(t)/t\, dt$ diverges if and only if $\int_1^{+\infty} f(t)^{\log q}\, dt$ diverges.

By the above lemma and as f is slowly varying, for every compact subset A of $\widehat{K} - \{\alpha, \alpha_*\}$, there exist positive constants c'_*, c''_* such that for every ξ in A,

- if $(r_n = [\gamma_n])_{n\in\mathbb{N}}$ is a sequence in R_0 with $D(r_n) \to +\infty$ as $n \to +\infty$ and $d_{C_0}(\xi, \gamma_n\alpha) \le c'_*\, f(D(r_n))\, e^{-D(r_n)}$ for every n big enough, then $h(\gamma_n\alpha) \to +\infty$ as $n \to +\infty$ and, for every n big enough,
$$|\xi - \gamma_n\alpha|_\infty^{\frac{1}{\log q}} \le e^{-g(-\log|\gamma_n\alpha - \gamma_n\alpha^*|_\infty^{\frac{1}{\log q}})}\, |\gamma_n\alpha - \gamma_n\alpha^*|^{\frac{1}{\log q}} ,$$
that is
$$|\xi - \gamma_n\alpha|_\infty \le \frac{\varphi(h(\gamma_n\alpha))}{h(\gamma_n\alpha)} ;$$
- conversely, if $(\gamma_n)_{n\in\mathbb{N}}$ is a sequence in Γ with $h(\gamma_n\alpha) \to +\infty$ as $n \to +\infty$ (in particular $\gamma_n \notin \Gamma_0$ for n big enough) and $|\xi - \gamma_n\alpha|_\infty \le \frac{\varphi(h(\gamma_n\alpha))}{h(\gamma_n\alpha)}$ for every n big enough, then with $r_n = [\gamma_n]$, we have $D(r_n) \to +\infty$ as $n \to +\infty$ and $d_{C_0}(\xi, \gamma_n\alpha) \le c''_*\, f(D(r_n))\, e^{-D(r_n)}$ for every n big enough.

Hence by Theorem 4.6, if $\int_1^{+\infty} \varphi(t)/t\, dt$ diverges, then for μ-almost every ξ in $\widehat{K}$, there exist a sequence $(\beta_n)_{n\in\mathbb{N}}$ in the congruence class of α in K_2, with $h(\beta_n) \to +\infty$ as $n \to +\infty$, such that $|\xi - \beta_n|_\infty \le \frac{\varphi(h(\beta_n))}{h(\beta_n)}$. This can be written as $\liminf \frac{\varphi(h(\beta))}{h(\beta)}\, |\xi - \beta|_\infty \le 1$, where the lower limit is taken over β in the congruence class of α with $h(\beta) \to +\infty$. Replacing φ by $\frac{1}{k}\varphi$ and letting k go to $+\infty$, this proves the divergence part of Theorem 1.6 in the introduction. The convergence part follows similarly. □

By taking $\varphi : t \mapsto t^{-s}$ in Theorem 1.6 with $s \ge 0$, the next result, which in particular says that almost every element of $\widehat{K}$ is badly approximable by quadratic irrational elements of $\widehat{K}$, follows immediatly.

Corollary 5.2 *For μ-almost every x in $\widehat{K}$,* $\liminf h(\beta)|x-\beta|_\infty = 0$*, and, for every $s>0$,* $\lim h(\beta)^{1+s}|x-\beta|_\infty = +\infty$*, where the lower limit and limit are taken over the quadratic irrational elements β in $\widehat{K}$, in any (resp. some) congruence class, with $h(\beta)\to+\infty$.* $\Box$

6 Approximating points

Let X be a proper CAT(-1) geodesic metric space. Let Γ be a non elementary discrete group of isometries of X, with finite critical exponent δ. In this section (which is a joint work with C. S. Aravinda), we will also apply our geometric avatar of the Borel-Cantelli Lemma, Theorem 3.1, to prove a Khintchine-type result for the approximation of a point by geodesic lines in X.

Let $x_0\in X$ be a base point. For every $C\geq 0$, a point ξ in $\partial_\infty X$ will be called a *C-strongly conical limit point* if there exist a geodesic line ρ with $\rho(+\infty)=\xi$ and a sequence $(\gamma_n)_{n\in\mathbb{N}}$ in Γ, such that $(\gamma_n x_0)_{n\in\mathbb{N}}$ converges to ξ, $d(\gamma_n x_0,\rho)\leq C$ and $d(\gamma_n x_0,\gamma_{n+1}x_0)\leq C$. Note that if ξ is a C-strongly conical limit point with respect to x_0, then ξ is a C'-strongly conical limit point with respect to any other base point x_0' for $C'=C+2\,d(x_0,x_0')$. And if ξ is a C-strongly conical limit point for the geodesic line ρ, then ξ is a $(C+\epsilon)$-strongly conical limit point with respect to any other geodesic line ρ' asymptotic to ρ, for every $\epsilon>0$.

Examples.

(1) If ξ is a fixed point of an hyperbolic element γ of Γ, then ξ is a C-strongly conical limit point with $C=\max\{d(x_0,A_\gamma),d(x_0,\gamma x_0)\}$, where A_γ is the translation axis of γ.

(2) If Γ is convex-cocompact, then there exists a constant $C\geq 0$ such that any limit point of Γ is a C-strongly conical limit point.

The following result is (a slight adaptation of) the well-known Sullivan's shadow lemma, see for instance [Bou1, page 93].

Lemma 6.1 *For every conformal density $(\mu_z)_{z\in X}$ of dimension δ for Γ, for every $C\geq 0$, there exists $c\geq 1$ such that for every C-strongly conical limit point ξ, for every $\epsilon\in\,]\,0,1]$,*

$$\frac{1}{c}\,\epsilon^\delta\leq\mu_{x_0}(B_{d_{x_0}}(\xi,\epsilon))\leq c\,\epsilon^\delta\ .\quad\Box$$

The following result is the main technical tool from which Theorem 6.3 in the introduction follows.

Theorem 6.2 *Let X be a* CAT(-1) *proper geodesic metric space. Let x,y be points in X. Let Γ be a non elementary discrete subgroup of isometries of X. Let $(\mu_z)_{z\in X}$ be a conformal density of dimension δ for Γ, for some δ in $]0,+\infty[$. Assume that there exist constants $c_0,C_0>0$ such that* $\operatorname{Card}\{\gamma\in\Gamma\ :\ d(x,\gamma y)\leq n\}\sim c_0\,e^{\delta n}$*, and that for every z in Γy except finitely many of them, there exists a geodesic ray ρ_z starting from x, passing through z and ending at a C_0-strongly conical limit point. Let $f:[\,0,+\infty[\,\to\,]\,0,+\infty[$ be a slowly varying map, with $f(t)$ converging to 0 as $t\to+\infty$. Let E_f be the set of points in $\partial_\infty X$ which belong to infinitely many balls $B_{d_x}\big(\rho_{\gamma y}(+\infty),f(d(x,\gamma y))\,e^{-d(x,\gamma y)}\big)$ for γ in Γ.*

[A] If $\int_1^\infty f(t)^\delta dt$ converges, then $\mu_x(E_f)=0$.

[B] If there exists a sequence $(t_n)_{n\in\mathbb{N}}$ *in* $\mathbb{R}$, *with* $t_n \to +\infty$ *as* $n \to +\infty$ *and* $t_{n+1} \geq t_n - \log f(t_n)$ *for every* n, *such that* $\sum_{n\in\mathbb{N}} f(t_n)^\delta$ *diverges, then* $\mu_x(E_f) > 0$.

Proof. Since Γ is discrete, the stabilizer Γ_y of y in Γ is finite, and there exists $c_1 \in \,]\,0,1]$ such that for every γ, γ' in Γ, if $d(\gamma y, \gamma' y) \leq 4\,c_1$, then $\gamma y = \gamma' y$, that is the class of γ and γ' in Γ/Γ_y are equal. It can be easily checked that for $[\gamma]$ in Γ/Γ_y, the objets $\rho_{\gamma y}$ and $d(x, \gamma y)$ are well defined.

Let $t'_0 > 0$ be big enough so that $f(t) \leq \frac{1}{e}$ for $t \geq t'_0$, and that for every γ in Γ such that $d(x, \gamma y) \geq t'_0$, the geodesic ray $\rho_{\gamma y}$ is defined. In case [A], define $t_0 = t'_0$ and by induction $t_{n+1} = t_n + 1$ for every n in $\mathbb{N}$. In case $[B]$, as $t_n \to +\infty$, we may assume, up to shifting the indices, that $t_0 \geq t'_0$.

For every n in $\mathbb{N}$, let

$$I_n = \{\gamma \in \Gamma/\Gamma_y \;:\; t_n \leq d(x, \gamma y) < t_n + c_1\}\;,$$

and $I = \bigcup_{n=0}^{\infty} I_n$. By the discreteness of Γ, the subsets I_n are finite, and pairwise disjoint since $c_1 \leq 1$, using also the hypothesis on the range of f in case [B].

Since f is slowly varying, there exists $c_2 \geq 1$ such that $f(y) \leq c_2 f(x)$ if $|y - x| \leq c_1$. Let $g = -\log f$, which is non negative on $[t'_0, +\infty[$, and $c_3 = \min\{c_1, \frac{1}{c_2}\}\, e^{-c_1}$, whicqh belongs to $]\,0,1]$. Note that the constants c_1, c_2, hence c_3, are unchanged if one replaces f by a scalar multiple of it. For every n in $\mathbb{N}$ and $\epsilon > 0$, define

$$f_1(n) = e^{\delta t_n},\;\; f_2(n) = c_3\, e^{-t_n},\;\; f_3(n) = c_3\, e^{-t_n - g(t_n)},\;\; f_4(n) = c_3^{-\delta},\;\; f_5(\epsilon) = \epsilon^\delta\;.$$

The series $\sum_{n\in\mathbb{N}} f_1(n) f_4(n) f_5(f_3(n))$ converges if and only if the series $\sum_{n\in\mathbb{N}} f(t_n)^\delta$ converges, which in case [A] is true if and only if the integral $\int_1^{+\infty} f^\delta$ converges, as f is slowly varying.

For every γ in I and $\epsilon > 0$, let $B_\gamma(\epsilon) = B_{d_x}(\rho_{\gamma y}(+\infty), \epsilon)$, which is measurable and non-decreasing in ϵ.

Let us prove that the finite measured space $(\partial_\infty X, \mu_x)$, the family $(B_\gamma(\epsilon))_{\gamma\in I,\, \epsilon>0}$, the finite-to-one map $I \to \mathbb{N}$ defined by $\gamma \mapsto n_\gamma = n$ if $\gamma \in I_n$, and the above maps f_1, f_2, f_3, f_4, f_5 satisfy the assumptions of Theorem 3.1.

Assumption (1) of Theorem 3.1 is satisfied as $g(t_n) \geq 0$ for every n. Assumption (2) is satisfied by the definition of f_4. Assumption (3) holds true with for instance $c' = 2, c'' = 2^\delta$. Assumption (4) follows by the hypothesis on the growth of the orbit Γy (counting elements of Γ/Γ_y rather than of Γ only divides the number by $\mathrm{Card}(\Gamma_y)$). Assumption (5) is satisfied by Lemma 6.1, as $f_2 \leq 1$ and f_4 is constant.

Let us prove Assumption (6) of Theorem 3.1. Given n in $\mathbb{N}$ and distinct α, β in I_n, assume by absurd that $B_\alpha(f_2(n))$ intersects $B_\beta(f_2(n))$ non trivially. By Lemma 2.1 (1), the ball $B_{d_x}(\rho_{\gamma y}(+\infty), c_1\, e^{-d(x,\gamma y)})$ is contained in ${}_xB(\gamma y, c_1)$ for every $\gamma \in I$. As $d(x, \alpha\, y) \leq t_n + c_1$ and $c_3 \leq c_1\, e^{-c_1}$, we have $f_2(n) \leq c_1\, e^{-d(x,\alpha\, y)}$, and similarly for β. Hence ${}_xB(\alpha\, y, c_1)$ intersects ${}_xB(\beta\, y, c_1)$ non trivially. There exists a geodesic ray ρ starting from x and passing at distance at most c_1 from both $\alpha\, y$ and $\beta\, y$. Let p, q be the closest point of $\alpha y, \beta y$ respectively on ρ, with (up to permuting α and β) $q \in [x, p]$. As $\alpha, \beta \in I_n$ and since the closest point maps do not increase distances, we have

$$d(\alpha\, y, \beta\, y) \leq d(p, q) + 2\, c_1 = d(x, p) - d(x, q) + 2\, c_1 \leq d(x, \alpha y) - d(x, \beta_y) + 3\, c_1 \leq 4\, c_1\;.$$

This contradicts the definition of c_1.

Finally, let us prove Assumption (7) of Theorem 3.1 under the hypotheses of Case [B]. For $m > n$, take (α, β) in $I_n \times I_m$, and assume that $B_\beta(f_3(m))$ intersects $B_\alpha(f_3(n))$ non trivially. Let w be a common point of these balls. Since $f_3 \leq f_2$, $t_m \geq t_{n+1}$ and $t_{n+1} \geq t_n + g(t_n)$, we have $f_3(m) \leq f_2(m) \leq f_2(n+1) \leq f_3(n)$. Then, for every z in $B_\beta(f_2(m))$, we have

$$\begin{aligned} d(z, \rho_{\alpha y}(+\infty)) &\leq d(z, \rho_{\beta y}(+\infty)) + d(\rho_{\beta y}(+\infty), w) + d(w, \rho_{\alpha y}(+\infty)) \\ &\leq f_2(m) + f_3(m) + f_3(n) \leq 3 f_3(n) \;. \end{aligned}$$

Therefore $B_\beta(f_2(m))$ is contained in $B_\alpha(3 f_3(n))$, which proves the claim.

Let E'_f be the subset of points of $\partial_\infty X$ which, as γ ranges over I, belong to infinitely many balls $B_\gamma(f_3(n_\gamma))$. As $0 \leq d(x, \gamma y) - t_n \leq c_1$ if $\gamma \in I_n$, and since $c_3 \leq \frac{1}{c_2} e^{-c_1}$, we have, for every γ in Γ,

$$B_\gamma(f_3(n_\gamma)) \subset B_{d_x}\big(\rho_{\gamma y}(+\infty), f(d(x, \gamma y))\, e^{-d(x,\gamma y)}\big) \subset B_\gamma\Big(\frac{c_2}{c_3} f_3(n_\gamma)\Big) \;. \tag{- 12 -}$$

In case [B], it follows from Theorem 3.1 that $\mu_x(E'_f) > 0$. By Equation (- 12 -), we have $\mu_x(E_f) > 0$.

In case [A], since the convergence or divergence of the integral $\int_1^{+\infty} f^\delta$ is unchanged if one replaces f by a scalar multiple of it, Theorem 3.1 implies that $\mu_x(E'_{\frac{c_2}{c_3} f}) = 0$. It follows similarly from Equation (- 12 -) that $\mu_x(E_f) = 0$. □

Given a complete Riemannian manifold M, recall (see [HP1]) that an element v in $T^1 M$, or its associated geodesic line ρ in M, is called *f-Liouville* at x_0 if there exists a sequence of times $(t_n)_{n \in \mathbb{N}}$ tending to $+\infty$ such that $d(\rho(t_n), x_0) \leq f(t_n)$ for every n. The following result is joint work with C. S. Aravinda.

Theorem 6.3 *Let M be a closed manifold with sectional curvature at most -1, let x_0 be a point in M, and let μ be the maximal entropy probability measure for the geodesic flow of M, with h its topological entropy. Let f be a slowly varying map.*

(1) If $f(t)$ and $e^{-t}/f(t)$ converge to 0 as $t \to +\infty$, if the integral $\int_u^\infty \frac{f(t)^h}{-\log f(t)}\, dt$ diverges (for some u big enough), then μ-almost every geodesic line is f-Liouville at x_0.

(2) If the sectional curvature of M satisfies $-a^2 \leq K \leq -1$, and if $\int_1^\infty f(t)^{\frac{h}{a}}\, dt$ converges, then μ-almost no geodesic line is f-Liouville at x_0.

Proof. Let $X \to M$ be a universal covering of M, with covering group Γ (which is non elementary), let x be a lift of x_0, and take $y = x$. For every z in $\Gamma y - \{y\}$, let ρ be the geodesic ray starting from x through z, which ends at a (uniformly) strongly conical limit point, as M is compact. Let $(\mu_z)_{z \in X}$ be the (unique up to positive scalar multiple) ergodic conformal density of dimension equal to the critical exponent δ of Γ (which is equal to h, as M is compact). Since M is compact, we have $\operatorname{Card}\{\gamma \in \Gamma \;:\; d(x, \gamma y) \leq n\} \sim c_0\, e^{\delta n}$ for some constant $c_0 > 0$. Hence the general hypotheses of Theorem 6.2 on $X, \Gamma, (\mu_z)_{z \in X}$ are satisfied.

Let us prove the first assertion of Theorem 6.3. Let $f_* : [0,+\infty[\to]0, \frac{1}{e}]$ be a slowly varying non increasing map such that $\frac{f_*(t)}{f(t)}$ tends to 0 as $t \to +\infty$ (so that in particular f_* converges to 0 at $+\infty$) and $\int_u^\infty \frac{f_*(t)^h}{-\log f_*(t)}\,dt$ still diverges. Fix $t_0 \geq 1$ such that $f_*(t_0) \leq \frac{1}{e}$ for $t \geq t_0$. Define by induction $t_{n+1} = t_n - \log f_*(t_n)$. In particular, the sequence $(t_n)_{n\in\mathbb{N}}$ converges to $+\infty$. As f_* is non increasing, we have

$$\int_{t_n}^{t_{n+1}} \frac{f_*(t)^\delta}{-\log f_*(t)}\,dt \leq \frac{f_*(t_n)^\delta}{-\log f_*(t_n)}\,(t_{n+1}-t_n) = f_*(t_n)^\delta\,.$$

Hence $\sum_{n\in\mathbb{N}} f_*(t_n)^\delta$ diverges. Therefore the hypotheses of Theorem 6.2 [B] are satisfied for f_*.

Denote by $\mathcal{S}$ the measurable set of elements ξ in $\partial_\infty X$ which belong to infinitely many visual balls $B_{d_x}\big(\rho_{\gamma x}(+\infty), f_*(d(x,\gamma x))\, e^{-d(x,\gamma x)}\big)$ as γ ranges over Γ. By Lemma 2.1 (1), this ball is contained in the shadow ${}_x\big(B(\gamma x, f_*(d(x,\gamma x)))\big)$. For every ξ in $\mathcal{S}$, let ρ_ξ be the geodesic ray starting from x and ending at ξ. As $f_*(t) \leq 1$ if $t \geq t_0$ and as f_* is slowly varying, there exist a constant $c_1 > 0$ and sequences $(s_n)_{n\in\mathbb{N}}$ in $[0,+\infty[$ converging to $+\infty$ and $(\gamma_n)_{n\in\mathbb{N}}$ in Γ such that $d(\rho_\xi(s_n), \gamma_n x) \leq c_1\, f_*(s_n)$ for every n. For every ξ in $\mathcal{S}$, let $\mathcal{S}_\xi$ be the set of elements v in T^1M such that the point at infinity of some geodesic line ρ_v, which is a lift by $X \to M$ of the geodesic line in M associated to v, is equal to ξ. Since X is CAT(-1), if ρ, ρ' are asymptotic geodesic rays, then there exists $c > 0$ and $\tau \in \mathbb{R}$ such that $d(\rho(t), \rho'(t+\tau)) \leq c\,e^{-t}$ for every $t \geq \max\{0, -\tau\}$. Since f is slowly varying, for every $v \in \mathcal{S}_\xi$, there exist hence a constant $c' > 0$ and sequences $(s'_n)_{n\in\mathbb{N}}$ in $[0,+\infty[$ converging to $+\infty$ and $(\gamma_n)_{n\in\mathbb{N}}$ such that

$$d(\rho_v(s'_n), \gamma_n x) \leq c'(f_*(s'_n) + e^{-s'_n}) \leq f(s'_n)\,,$$

for n big enough. Hence every $v \in \mathcal{S}_\xi$ is f-Liouville, for every $\xi \in \mathcal{S}$. By Theorem 6.2 [B], the set $\mathcal{S}$ has positive measure for μ_x. Hence the set $\mathcal{S}' = \bigcup_{\xi\in\mathcal{S}} \mathcal{S}_\xi$ (which is measurable) has positive measure for μ, by Lemma 2.4. As $\mathcal{S}'$ is invariant under the geodesic flow by construction, and by ergodicity, it has full measure. This proves the result.

Let us now prove the assertion (2) of Theorem 6.3. As f is slowly varying, the convergence of the integral $\int_1^\infty f^{\frac{\delta}{a}}$ implies that f converges to 0 as $t \to +\infty$. For every v in T^1M, let ρ_v be a lift by $X \to M$ of the geodesic line in M associated to v, let $\xi_v = \rho_v(+\infty)$ and let ρ'_v be the geodesic ray from x to ξ_v. If v is f-Liouville at x_0, then there exist sequences $(s_n)_{n\in\mathbb{N}}$ in $[0,+\infty[$ converging to $+\infty$ and $(\gamma_n)_{n\in\mathbb{N}}$ in Γ such that $d(\rho_\xi(s_n), \gamma_n x) \leq f(s_n)$ for every n. Since f is slowly varying and since ρ'_v and ρ_v are asymptotic, as above, there exist k in $\mathbb{N} - \{0\}$, a sequence $(s'_n)_{n\in\mathbb{N}}$ in $[0,+\infty[$ converging to $+\infty$, and $(\gamma_n)_{n\in\mathbb{N}}$ in Γ, such that $d(\rho'_v(s'_n), \gamma_n x) \leq k(f(s'_n) + e^{-s'_n})$ for every n. In particular, as $f(t) \leq 1$ if t is big enough, if v is f-Liouville, then there exists k' in $\mathbb{N} - \{0\}$ such that ξ_v belongs to infinitely many shadows ${}_x\Big(B\big(\gamma x, k'(f(d(x,\gamma x)) + e^{-d(x,\gamma x)})\big)\Big)$ for $\gamma \in \Gamma$. By Lemma 2.1 (2), this shadow is contained, except for finitely many $\gamma \in \Gamma$, in the ball

$$\mathcal{B}_{\gamma,k''} = B_{d_x}\Big(\rho_{\gamma x}(+\infty), k''\big(f(d(x,\gamma x)) + e^{-d(x,\gamma x)}\big)^{\frac{1}{a}}\, e^{-d(x,\gamma x)}\Big)\,,$$

for some positive integer k''. If $u, v, w > 0$, recall that $(u+v)^w \leq 2^w(u^w + v^w)$. Hence since $\int_1^\infty f^{\frac{\delta}{a}}$ converges, the integral $\int_1^\infty \left(k''(f(t)+e^{-t})^{\frac{1}{a}}\right)^\delta \, dt$ also converges. The map $t \mapsto k''(f(t)+e^{-t})$ is slowly varying. By Theorem 6.2 [A], the measure of the set of points in $\partial_\infty X$ which belong to infinitely many balls $\mathcal{B}_{\gamma,k''}$, as γ ranges over Γ, has measure 0 for μ_x. By Lemma 2.4, and since a countable union of measure zero subsets is a measure zero subset, the result follows. $\square$

For every $\alpha > 0$, let $f_\alpha : t \mapsto \frac{1}{(2+t)^\alpha}$, which is slowly varying, with $f(t)$ and $e^{-t}/f(t)$ converging to 0 as $t \to +\infty$. For every $h > 0$, the integral $\int_1^\infty \frac{f_\alpha^h}{-\log f_\alpha}$ diverges if and only if $\alpha \leq \frac{1}{h}$ and the integral $\int_1^\infty f_\alpha^h$ converges if and only if $\alpha > \frac{1}{h}$. By applying Theorem 6.3 with M having constant curvature -1, so that $h = n-1$, with $f = f_{\frac{1}{h} \pm \frac{1}{n}}$ where $n \to +\infty$, Theorem 1.7 of the introduction follows, in the standard way one deduces a logarithm law-type theorem from a Khintchine-type theorem.

References

[Arz] G. Arzhantseva, *On quasi-convex subgroups of word hyperbolic groups*, Geom. Dedicata **87** (2001) 191–208.

[Bea] A. Beardon, *The geometry of discrete groups*, Springer-Verlag, 1983.

[BV] V. Beresnevich, S. Velani, *Ubiquity and a general logarithm law for geodesics*, preprint Univ. York, July 2007. [`arXiv:0707.1225`]

[Bou] N. Bourbaki, *Topologie générale*, chap. 1 à 4, Hermann, Paris, 1971.

[Bou1] M. Bourdon, *Structure conforme au bord et flot géodésique d'un CAT(-1) espace*, L'Ens. Math. **41** (1995) 63–102.

[Bou2] M. Bourdon, *Immeubles hyperboliques, dimension conforme et rigidité de Mostow*, GAFA **7** (1997) 245–268.

[Fur] H. Furusawa, *The exponent of convergence of Poincaré series of combination groups*, Tohoku Math. J. **43** (1991) 1–7.

[BH] M.R. Bridson, A. Haefliger, *Metric spaces with non-positive curvature*, Grund. math. Wiss. **319**, Springer Verlag (1998).

[BP] A. Broise, F. Paulin, *Dynamique sur le rayon modulaire et fractions continues en caractéristique p*, to appear in Journal of the London Mathematical Society. [`math.GR/0511442`]

[DMPV] M. M. Dodson, M. V. Meliàn, D. Pestana, S. L. Velani, *Patterson measure and ubiquity*, Ann. Acad. Sci. Fenn. **20** (1995) 37–60.

[GP] D. Gaboriau, F. Paulin, *Sur les immeubles hyperboliques*, Geom. Dedi. **88** (2001) 153–197.

[GH] E. Ghys, P. de la Harpe, eds. *Sur les groupes hyperboliques d'après Mikhael Gromov*, Prog. in Math. **83**, Birkhäuser 1990.

[HaP] F. Haglund, F. Paulin, *Constructions arborescentes d'immeubles*, Math. Ann. **325** (2003) 137–164.

[Ham] U. Hamenstädt, *A new description of the Bowen-Margulis measure*, Erg. Theo. Dyn. Sys. **9** (1989) 455–464.

[HPP] S. Hersonsky, J. Parkonnen, F. Paulin, *Spiraling spectra of geodesics in negative curvature*, in preparation.

[HP1] S. Hersonsky, F. Paulin, *On the rigidity of discrete isometry groups of negatively curved spaces*, Comm. Math. Helv. **72** (1997) 349–388.

[HP2] S. Hersonsky, F. Paulin, *Hausdorff dimension of diophantine geodesics in negatively curved manifolds*, J. reine angew. Math. **539** (2001) 29–43.

[HP3] S. Hersonsky, F. Paulin, *Diophantine approximation for negatively curved manifolds, I*, Math. Zeit. **241** (2002) 181–226.

[HP4] S. Hersonsky, F. Paulin, *Counting orbit points in coverings of negatively curved manifolds and Hausdorff dimension of cusp excursions*, Erg. Theo. Dyn. Sys. **24** (2004) 1–22.

[HP5] S. Hersonsky, F. Paulin, *A logarithm law for automorphism groups of trees*, Archiv Math. **88** (2007) 97-108.

[HS] M. Hindry, J. Silverman, *Diophantine geometry : an introduction*, GTM **201**, Springer Verlag, 2000.

[Kai] V.A. Kaimanovich, *Bowen-Margulis and Patterson measures on negatively curved compact manifolds.* in “ Dynamical systems and related topics” (Nagoya, 1990), 223–232, Adv. Ser. Dyn. Syst., **9**, World Sci. Pub., 1991.

[Kap] M. Kapovich, *Representations of polygons of finite groups*, Geom. Topol. **9** (2005) 1915–1951

[KS] I. Kapovich, H. Short, *Greenberg's theorem for quasiconvex subgroups of word hyperbolic groups*, Canadian J. Math., **48** (1996) 1224–1244.

[KM] D. Kleinbock, G. Margulis, *Flows on homogeneous spaces and Diophantine approximation on manifolds*, Ann. Math. **148** (1998) 339–360.

[Kni] G. Knieper, *Spherical means on compact Riemannian manifolds of negative curvature*, Diff. Geom. App. **4** (1994) 361–390.

[Las] A. Lasjaunias, *A survey of Diophantine approximation in fields of power series*, Monat. Math. **130** (2000) 211–229.

[Leu] E. Leuzinger, *Kazhdan's property (T), L^2-spectrum and isoperimetric inequalities for locally symmetric spaces*, Comment. Math. Helv. **78** (2003) 116–133.

[Man] A. Manning, *Topological entropy for geodesic flows*, Ann. Math. **110** (1979) 567–573.

[PP] J. Parkonnen, F. Paulin, *Prescribing the behaviour of geodesics in negative curvature*, preprint Univ. Jyvaskyla, 2007. [`arXiv:0706.2579`]

[Pau] F. Paulin, *Groupe modulaire, fractions continues et approximation diophantienne en caractéristique p*, Geom. Dedi. **95** (2002) 65–85.

[Rob] T. Roblin, *Ergodicité et équidistribution en courbure négative*, Mémoires Soc. Math. France, **95** (2003).

[Sch] W. Schmidt, *On continued fractions and Diophantine approximation in power series fields*, Acta Arith. **XCV** (2000) 139–166.

[Ser] J.-P. Serre, *Arbres, amalgames, SL_2*, Astérisque **46**, Soc. Math. France, 1983.

[Spr] V.G. Sprindzŭk, *Mahler's problem in metric number theory*, Transl. Math. Mono **25**, Amer. Math. Soc., 1969.

[SV] B. Stratmann, S. L. Velani, *The Patterson measure for geometrically finite groups with parabolic elements, new and old*, Proc. London Math. Soc. **71** (1995) 197–220.

[Sul] D. Sullivan, *Disjoint spheres, approximation by imaginary quadratic numbers, and the logarithm law for geodesics*, Acta Math. **149** (1982) 215–237.

[SS] P. Susskind, G. A. Swarup, *Limit sets of geometrically finite hyperbolic groups*, Amer. J. Math. **114** (1992) 233–250.

University of Georgia
Department of Mathematics
Athens, GA 30602, USA
e-mail: saarh@math.uga.edu

and

Ben Gurion University
Department of Mathematics
Bear Sheva, Israel
e-mail: saarh@cs.bgn.ac.il

Département de Mathématique et Applications, UMR 8553 CNRS
École Normale Supérieure, 45 rue d'Ulm
75230 PARIS Cedex 05, FRANCE
e-mail: Frederic.Paulin@ens.fr